

www.pmc.edu.bd

ISSN-L:2958-7794

Volume 2, Issue 1, May 2023

PMJ



Pabna Medical College Journal

Editorial

- Artificial Intelligence and Medical Science: Present Perspective 1
Ahmmad Taous

Original Articles

- Percutaneous Dilatational Tracheostomy: an Observational Study at Birat Medical College and Teaching Hospital, Kathmandu, Nepal 3
Shah RK, Ullah MS, Koirala B, Rajbansi LK, Hasan DM, Arjyal B
- The “Lelli’s Test”: A Screening Clinical Test for the Diagnosis of Anterior Cruciate Ligament Injury 7
Ali MM, Naharin NE, Hossain SME, Manjur M, Islam MN, Masur MAA, Khatun MF
- Factors Related to Respiratory Distress Syndrome in Neonates 11
Akhund M, Hamid S, Hossain I, Alam UK, Noor IN
- Comparative Study of Neck Swelling by Clinical, Cytological and Histopathological Examination – A Study of 50 Cases 18
Ali MI, Huq MM, Rahman SH, Tarafder KH
- Visual Acuity of Children of Darisharifpur Village, Pabna 27
Mridha MZEA

PABNA MEDICAL COLLEGE JOURNAL (PMJ)

Official Journal of Pabna Medical College, Bangladesh

EDITORIAL BOARD

Chairperson

Prof. Dr. Md. Obaidullah Ibne Ali

Executive Editor

Prof. Dr. Ahmmad Taous

Editor

Prof. Dr. Abu Mohd. Shafiqul Hasan

Assistant Editors

Dr. Biplob Kumar Saha

Dr. Md. Sherajul Islam

Dr. Mohammad Habibullah

Members

Dr. Md. Mahbubul Alam

Dr. Md. Pervez Amin

Dr. Nargis Sultana

Dr. Md. Ruhul Quddus

Dr. Sougata Mitra

Dr. Md. Mizanur Rahman

Dr. Kh. Mehedy Ibnay Mostofa

Dr. Shamima Yeasmin

Advisors

Prof. Dr.Md. Reazul Haque

Prof. Dr. Iftekhar Mahmood

Prof. Dr.Md. Rayhanur Rahman

Dr. Shaheen Ferdous

Dr. Kazi Mohiuddin Ahmed

Office : Room No- 223, First floor, Pabna Medical College, Pabna
Phone- +880 1717033773, +880 258882083
Email: atlincoln71@gmail.com, amsshahan@gmail.com

Published by : Teachers Association, Pabna Medical College, Pabna, Bangladesh
Phone: +88 01717033773, Email: pmc@dghs.gov.bd

Printed at : Asian Colour Printing, 130, DIT Extension Road, Fakirerpool, Dhaka-1000, Bangladesh.
Tel: +88 02 58313186, +88 01835180135, E-mail: asianclr@gmail.com Web: asiancolourprinting.com

PABNA MEDICAL COLLEGE JOURNAL (PMJ)

Official Journal of Pabna Medical College, Bangladesh

PEER REVIEW PANEL

Professor Dr. Md. Mujibur Rahman

Professor of Medicine
BSMMU, Dhaka
Phone: +8801711525406
Email: mmrahman61@gmail.com

Professor Dr. Md. Rayhanur Rahman

Professor of Surgery
Pabna Medical College, Pabna
Phone: +88 01711 284 791
Email: md.rayhanur@yahoo.com

Professor Dr. Md. Bulbul Hasan

Professor of Microbiology
Naogaon Medical College, Naogaon
Phone: +8801740 980 319
Email: hasanmdbulbul@gmail.com

Professor Dr. Sheikh Nurul Fattah Rumi

Professor of ENT
Dhaka Medical College, Dhaka
Phone: +880 1711452381
Email: rumi17dr@gmail.com

Professor Dr. Md. Obaidullah Ibne Ali

Professor of Physiology
Pabna Medical College, Pabna
Phone: +8801711 014 82
Email: dr.bachchu1966@gmail.com

Professor Dr. Ahmmad Taous

Professor of ENT
Pabna Medical College, Pabna
Phone: +88017117033773
Email: atlincoln71@gmail.com

Professor Dr. Md. Khalilur Rahman

Professor of Medicine
Rajshahi Medical College, Rajshahi
Phone: +8801711302261
Email: drkhalil64@yahoo.com

Professor Dr. Abu Mohd. Shafiqul Hasan

Professor of Medicine
Pabna Medical College, Pabna
Phone: +880 1711 581 685
Email: amsshasan@gmail.com

Professor Dr. Md. Habibullah Sarker

Professor of Surgery
Rajshahi Medical College, Rajshahi
Phone: +8801711626953
Email: habibsarkar66@gmail.com

Professor Dr. Md. Shafiqul Bari

Professor of Medicine
Dhaka Medical College, Dhaka
Phone: +8801726948138
Email: drsbari69@yahoo.com

Pabna Medical College Journal (PMJ)

Official Journal of Pabna Medical College, Bangladesh

ABOUT US

Focus and Scope

Pabna Medical College Journal is a biannual, open access, double-blinded peer-reviewed journal. It publishes original articles, editorials and case reports based on Community, Clinical and hospital based laboratory work, field work, clinical trials and various other studies by scientific means related to the disciplines of medical and health sciences. It also accepts review articles, meta-analysis, conference proceedings, news related to medical science and letters to the editor.

Peer Review Process

Each manuscript is initially collected and screened by the editors. If it is considered for review process, then it is sent to members of preselected reviewer panel. The peer reviewers suggest editing, correction or rewriting the manuscript. After completion of the review process, the manuscript is selected for upcoming issue.

Authorship policy

The authors and Co-authors should have major contribution in research or study. They will take the public responsibility of the research procedure, findings or recommendation.

Selection of Reviewers

Reviewer selection is the critical and important part to the publication process. The choice of selection is based on several factors, including a) expertise, b) reputation, c) capability of giving recommendations, d) giving enough time to review and e) our own previous experience of a reviewer's characteristics. For example, the reviewer who is slow, careless, or do not provide definite comment is avoided.

Editorial Decision

The editor takes the final decision based on the reviewers' comment. The decision may be any one of the followings:

- a) Accept for Publication, with or without any minor editorial revision
- b) Inform the corresponding author to Revise the Manuscript to address the specific concerns before a final decision is reached. Submission of revised manuscript does not ensure the acceptance
- c) Reject the manuscript and suggests for rewriting and Resubmission.

Ethical issue

When reporting experiments on human subjects, authors have to write whether the procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000. When reporting experiments on animals, authors should indicate whether the institutional and national guide for the care and use of laboratory animals was followed. The work will be rejected if there is lack of evidence.

There is an Ethical Committee of Pabna Medical College to guide the authors to follow the rights of subjects of study or on experiment.

Conflict of interest

The credibility of published article depends in part on how well conflict of interest is handled during writing, peer review, and editorial decision making. Conflict of interest exists when an author, reviewer, or editor has financial or personal relationships that inappropriately influence (bias) his or her actions (such relationships are also known as dual commitments, competing interests, or competing loyalties). All authors of the submitted manuscript are requested to disclose any actual or potential conflict of interest including any financial, personal or other relationships with other people or organizations within two years of beginning the submitted work that could

inappropriately influence, or be perceived to influence, their work. If these conflicts occur, they have to report as well.

Confidentiality

Editor, editorial board members, authors and reviewers keep confidential of all details of the editorial and peer review process on submitted manuscripts. Even identities of the reviewer are not disclosed to the corresponding author. In addition, reviewers maintain the confidentiality of the manuscript.

Publication Frequency

The journal is publishing 2 issues (may and November) in a year from January 2023.

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

Publisher

Teachers Association, Pabna Medical College, Pabna, Bangladesh

Pabna Medical College Journal (PMJ)

Official Journal of Pabna Medical College, Pabna, Bangladesh

GUIDELINES FOR AUTHORS

The Authors may submit articles electronically round the year following the guideline of journal. The editors reserve the right to reject articles without sending them out for review. Submitted articles should not have been previously published or be currently under consideration for publication elsewhere. Briefs and research notes are not published in this journal. Submitted articles must be within the scope of the journal. All articles go through a peer review policy.

Authors must acknowledge that they have read and agree to the content of the submitted article with an acceptable standard of good English grammar and usage.

Article Processing Charges (APC)

As an open access journal with no subscription charges, a fee (Article Processing Charge, APC) is payable by the author or research funder to cover the costs associated with publication.

Types of Articles

Authors must ensure that while submitting the article, it should fall under the appropriate category from the list of options:

1. **Original Research** – Original research articles should describe confirmed findings and experimental methods should be given with sufficient details for others to verify the work. The length of an original article should describe and interpret the work clearly.
2. **Article Review** – Review articles, often known as literature reviews or secondary sources, synthesize or assess primary source research. In general, they summarize the status of research in a certain area. Review articles usually focus on presenting and discussing recent developments in a field.
3. **Case Study** – A case study is a detailed study of a specific area that involves researching a problem, often using an experiment or survey,

and providing potential solutions from that research. It is an in-depth look at one research problem, looking at it from different angles and aspects.

Submission Checklist:

Author (s) can use this list to carry out a final check of article submission before sending it to the journal for peer review process. Authors should ensure to have the following items in the article which should be available:

1. **Authorship:**

Authorship provides credit and bears public responsibility for a researcher's contribution to a study.

 - a. *Authors Contacts:* Authors should have been designated with contact details: Full postal address, E-mail address, and ORCID or Scopus ID or provide link i.e. Google scholar etc in his/her corresponding address.
 - b. *Corresponding Authors (CAs):* CAs is authors and act as a communicator between authors and the journal. CAs provides the basic requirements of articles to the Journal. Preferably CAs will be the principal author. This responsibility incorporates answering any future queries about methodology and materials. Ensure that the e-mail address is given, and the contact details are kept up to date by the corresponding author.
2. **Basic Requirements for the Article- A Brief Summary:**
 - a. *Words Count:* Not more than 4000 words (excluding Abstract, Methods, Tables, References, and figure legends).
 - b. *Page Range:* Not more than 12 pages.
 - c. *Submitting File Size:* Not more than 5MB (.doc/.docx/ PDF).

- d. *Title*: Not more than 50 words.
- e. *Abstract*: Minimum of 150 words and maximum of 250 words.
- f. *Keywords*: Minimum of 3 and maximum of 5 appropriate keywords.
- g. *Font Name and Font Size*: Times New Roman, Font Size 12, single or double column with 1.5 line spacing and justified.
- h. *Figures*: Not more than 5 figures. The figures should be in 300 dpi or more.
- i. *Tables*: Not more than 8 tables. The length of the table should not be more than 01 page.
- j. *Figure Legends*: Not more than 150 words per figure.
- k. *Graphical Abstracts / Highlights files*: Optional.
- l. *Supplemental files*: Optional
- m. *Spell and Grammar*: Compulsory.
- n. *Abbreviations*: Not more than 10 abbreviations. Ensure consistency of abbreviations throughout the article.
- o. *Declaration*: Authors are required to include a declaration of accountability in the article, counting review-type articles that stipulate the involvement of each author. The level of detail differs; some subjects yield articles that consist of isolated efforts that are easily voiced in detail, while other areas function as group efforts at all stages. It should be after the conclusion and before the references.
 - i. *Conflicts of Interest/ Competing Interests*: The article should not be under Conflict of Interest.
 - ii. *Funding/ Grants/ Financial Support*: Authors should detail if any financial support received.
 - iii. *Ethical Approval and Consent to Participate*: The authors must submit a statement that the study does not require ethical approval and consent to participate with evidence.
 - iv. *Availability of Data and Material/ Data Access Statement*: The Authors should mention source of research data and data access terms and conditions in the article.
- v. *Authors Contributions*: Authors should describe the role of each author if there is more than one author. See authorship for more details.
- p. *References*: Not more than 60 references.
- q. *Referee suggestions*: To be bestowed based on journal requirements.
- r. *Footnote*: Footnote should not be employed.

Article Format/Structure:

Article text file (ms word/pdf) should start with a title page that shows author affiliations and contact information, identifying the corresponding author with an asterisk. For the main body of the text, there are no specific requirements. It can be organized in a way that best suits the research.

Do not use 'all capital' letter in headings or subheadings, titles or subtitles.

However, the following structure will be suitable in many cases:

1. **Abstract**: Do not incorporate any references in Abstract. Verify they serve both as a general introduction to the topic and as a brief, non-technical summary of the main results and their implications.
2. **Keywords/index terms**: There should be appropriate keywords.
3. **Introduction**: Articulate the objectives of the work and bestow an adequate background, avoiding a detailed literature survey or a summary of the results.
4. **Materials and Methods**: Specify sufficient particulars to favour the work to be reproduced by an independent researcher. Methods that are already published should be illustrated and indicated by a reference. If quoting directly from a previously published method, use quotation marks and cite the source. Any modifications to existing methods should be proclaimed.
5. **Results (with subheadings if applicable)**: Results should be clear and concise.
6. **Discussion (without subheadings)**: This should explore the significance of the results of

the work without repetition. A combined Results and Discussion section is often appropriate. Avoid extensive citations and discussion of published literature.

7. **Conclusion:** The main conclusions of the study should be presented in a short Conclusions section, which may stand alone or form a subsection of a Discussion or Results and Discussion section.
8. **Acknowledgement:** Individuals who imparted cooperation during the research (e.g. providing language help, writing assistance, or proof reading the article, etc.) should be listed here.
9. **Figures:** Serial number of figure should be written in English numerical. Keep text in the illustrations themselves to a minimum but clearly explain all symbols and abbreviations used.
10. **Tables:** All tables should be editable as text and not as images. Serial number of tables should be given in Roman numerical. Usually does not require any description of tables, avoid duplicity of the data presented in them described elsewhere in the article.
11. **References:** Ensure that all references mentioned in the reference list are cited in the text and vice versa.

Abstract:

A concise and factual abstract is mandatory. The abstract should be structured, briefly state the intention of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. It does not require references. Also uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Keywords:

Promptly after the abstract, bestow a maximum of 6 keywords, using American spelling and avoid general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field will be eligible. These keywords will be used for indexing purposes.

Cover Letter:

The submission must also comprise a cover letter. In the cover letter, author should incorporate:

1. The affiliation, and contact information of the corresponding author.
2. The Affiliation of all co-authors.
3. Declare that the submitted article was not published before in any other journal.

Methods:

If author is reporting experiments on live vertebrates (or higher invertebrates), humans or human samples, author must comprise a statement of ethical approval in the Methods

References:

Edit references are not copied. Therefore, it is essential that the author formats them accurately, as they will be linked electronically to external databases where feasible. So, when formatting references, ensure that:

1. Run sequentially (and are always numerical).
2. Only have one publication associated to each number.
3. Only comprise articles or datasets that have been published or accepted by a named publication, recognized preprint server or data repository (if author include any preprints of accepted papers in reference list, ensure to submit them with the article).
4. Incorporate published conference abstracts and numbered patents if authors wish.

In reference list, author should:

1. Incorporate all authors unless there are six or more, in that case 6 author should be mentioned, followed by 'et al.'
2. Classify authors by last name first, followed by a comma and initials (followed by full stops) of given names.
3. Adopt Roman text for Article and dataset titles, with only the first word of the title having an initial capital and written exactly as it appears in the work cited ending with a full stop.

4. Apply Italics for book titles, giving all words in the title an initial capital.
5. Apply Italics for journal and data repository names, abbreviating them according to common usage (with full stops).
6. Contribute the full-page range (or article number) where appropriate.

Examples:**Published Article in journal**

Burks RT, Morgan J. Anatomy of the lateral ankle ligaments. *Am J Sports Med* 1994; 22 (1): 53-67

Books:

Alaszewski A. *COVID-19 and Risk: Policy Making in a Global Pandemic*. Policy Press; 2021.

Book chapter:

Wynick D, Bloom SR. Islet cell tumors. In: Grossman A. *Clinical Endocrinology*. Oxford: Blackwell Scientific Publications, 1992: 502-511.

Personal Author:

Adam RD, Victor M. *Principles of Neurology*. 5th Ed. New York: McGraw-Hill, 1993.

Editor(s) or Compiler(s):

Diener HC, Wilkinson M, editors. *Drug-induced headache*. New York: Springer-Verlag, 1988.

Organization as author:

Virginia Law Foundation. *The Medical and legal implications of AIDS*. Charlottesville: The Foundation, 1987.

Online Material:

Babichev SA, Ries, J. & Lvovsky, A. I. Quantum scissors: teleportation of single- mode optical states by means of a nonlocal single photon. Preprint at <https://arxiv.org/abs/quant-ph/0208066> (2002).

Website citation:

Pelc C. Prescription steroids affect brain structure, study finds. Published September 6, 2022. Accessed September 8, 2022. <https://www.medicalnewstoday.com/articles/prescription-steroids-affect-brain-structure-study-finds>.

Acknowledgements:

Aggregate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who contributed cooperation during the research (e.g. providing language help, writing assistance or proof reading the article etc.).

Author Contributions:

The author must supply an Author Contribution Statement as described in the Author Responsibilities section.

Please be aware:

1. The author's name which author gives as the corresponding author will be the main contact amid the review process and should not be changed.
2. The information author bestows in the submission system will be used as the source of truth when his/her article is published.

Pabna Medical College Journal (PMJ)

Official Journal of Pabna Medical College, Bangladesh

Publication and Data Policy

Preparation and submission of Manuscript:

Before submission, the authors are requested to strictly follow the guide lines below for submission of manuscript. Manuscripts prepared following the **Uniform Requirements for manuscripts submitted to biomedical Journals** is acceptable to this journal for publishing. Authors are advised to visit the web site www.icmje.org for this purpose. In addition, authors are requested to watch the format and style of the article published in this journal. The following documents with manuscripts are to be submitted for publication. A covering letter addressed to the Editor of the journal. A Title page. Abstract and key words in the first page followed by the text, prepared in the format of IMRAD. Authors must submit 2 hard copies of all documents and one copy in electronic form preferably written in a CD with adequate labeling. As an alternative option, submission through email with file attachment of all documents is welcome.

Covering letter:

All authors must sign after seeing the manuscript with the statement that they are the only author. The corresponding author should mention the contribution of each author to the work. It should contain a declaration that this manuscript has not been submitted elsewhere or not under consideration in any journal. It should clearly indicate the publication type (Original/Review/case report/letter etc.) Authors are requested to submit new and revised manuscript and correspondence to: Editor Pabna Medical College Journal Email: pmj12345@gmail.com, Phone no. +880 17171033773, +8801711581685

Title page:

It should contain title of the article followed by Name and affiliation of each byline author, the center/centers to which the work is attributed, name, address, telephone number, email address of corresponding author, declaration of conflict of interest, disclaim if any, disclosure for financial or other support, number of tables, number of figures given and a short running title and lastly word count of abstract.

Manuscript Organization:

It should be typed double spaced throughout with Justified and 2.5 cm margins. Font type is Times New Roman with size 12 and printed on a good quality A4 size weight 80 gm of paper on one side. Manuscript should have uniform style, correct journal format, carefully proofread for grammar, spelling and punctuation.

Manuscript format:

In general, original article should be divided into following sections: Title page, Abstract, Text, Acknowledgement, References. At the end, Tables with titles and foot notes, alternatively graph with title and Illustrations with legends. Each of the sections is to start on a separate page. Pages should be numbered consecutively beginning from the abstract.

Abstract page:

It should start with appropriate title of the article and structured with headings (Background, Objectives, Methods with statistical analysis, Result & Conclusion. Authors name should not be given at this page. It should be preferably within about 250 words. Avoid abbreviations in the title and abstract except standard abbreviation. A non structured abstract is suggested for review article and case report. It should be followed by 3 to 5 Key words.

Text:

Text should be arranged into following sections: Introduction, Methods, Results, Discussions, Acknowledgement, References.

Introduction: It will include statement of the problem with a short discussion of its importance and significance, review of the literature related to the problem with pertinent reference and lastly Objectives/hypothesis/ benefits expected stated in 1-2 paragraph.

Methods: In this section study type, place and time, description of study subjects and grouping, sampling, selection criteria, approval of the study involving human subjects by ethical review committee and description of the ethical aspects in such study,

description of study variable, description of procedure, methods, apparatus, drugs or chemicals as applicable should be given. Description of statistical procedure with enough detail to enable a knowledgeable reader with access to the original data to verify the reported results should be provided. For animal experimental study, animal ethical aspects should also be mentioned.

Results: Present result in logical sequence in text, table and illustration with most important finding first. Describe without comment, restrict number of table and figure needed to support assessment of paper. Do not duplicate data in table and figure.

Discussion: In this section, describe only the key findings and their comparison to other researchers, their interpretation, probable explanation from literature if relevant and the message of the study.

Table: Simple self explanatory with brief title, not duplicate in text. Each table should be numbered in **Romans** and printed in separate page. Do not use internal horizontal and vertical rules. Uses of many tables are not encouraged. It should be mentioned in result section.

Illustration: All illustrations must be numbered consecutively in **Arabic numerals** as they appear in the text. Submit Print photograph of each illustration along with its electronic file. Figure number, title of manuscript, name of the corresponding author and arrow indicating top should be written on a sticky label affixed on the back of each photograph. Scanned picture, graph, chart with title and figure number should be printed on separate page and its original data presentation file should be inserted in the CD along with text.

Legend: Must be typed in a separate sheet of paper. Photomicrograph should indicate the magnification, internal scale and the method of staining. All drugs should be mentioned in their generic form. The commercial name may be used in parenthesis.

Acknowledgement: Individuals, institution, sponsor, organization or bodies can be acknowledged in the article for their contribution or financial or any form of assistance to the work.

References: For reference, use author number style (Vancouver) which is based on an ANSI standard adapted by the National Library of Medicine (NLM). References should be numbered consecutively in the order in which they are first mentioned in the text. Identify references in the text, tables and legends by Arabic numerals in **superscript**. All

citations to electronic references should be presented in numbered references following the text. The titles of the journals should be abbreviated according to the style used in Index Medicus. Write names all authors of the article. **The reference list is not usually checked by the editorial staff or reviewer. It is the total responsibility of author to provide accurate information.**

Examples:

Standard journal article:

Khalil M, Chowdhury MAI, Rahman H, Mannan S, Sultana SZ, Rahman MM et.al. Splenic Mass and its relation to age, sex and height of the individual in Bangladeshi Peoples. J Bangladesh Soc Physiol 2008;3(1):71-78.

Journal article with organization as author:

Name of organization full stop title of the article full stop abbreviated journal title comma year and month of publication semicolon volume colon page number.

American diabetes Association. diabetes update. Nursing, 2003 Nov; Suppl: 19-20.

Journal article with multiple organization as author:

Name of the 1st organization semicolon Name of the 2nd organization semicolon Name of the 3rd organization full stop title of the article full stop abbreviated journal title full stop year and month of publication semicolon volume colon page number.

American Dietetic association; Dietitians of Canada. Position of Dietetic association and Dietitians of Canada: nutrition and women's health. J Am Diet Assoc. 2004 Jun; 104(6):984-1001.

Journal article with Governmental body as author:

Name of the bodies with intervening semicolon end with full stop topic colon title of article full stop Abbreviated journal title full stop year and month of publication semicolon volume colon page number.

National Institute on Drug Abuse (US); Caribbean Epidemiology Centre; Pan American Health Organization; World Health Organization. Building a collaborative research agenda: drug abuse and HIV/AIDS in the Caribbean 2002-2004. West Indian Med J. 2004 Nov; 53 Suppl 4:1-78

Standard book with initials for authors:

Name of the authors full stop title of the book full stop edition full stop place of publication colon name of publisher semicolon copy right year full stop page number p full stop

Eyre HJ,LangeDP,MorrisLB.Informed decisions: the complete book of cancer diagnosis,treatment and recovery.2nd ed. Atlanta: American Cancer Society; 2002.768p.

Contributed chapter of a book:

Name of the authors of chapter full stop title of the chapter full stop In colon name of editors/editor coma editors/ edito full stop title of the book full stop place of publication colon name of publisher semicolon copy right year full stop page ranges.

Rojko JL, Hardy WD. Feline leukemia virus and other retroviruses. In:SherdingRG,editor.Thecat: diseases and clinical management.New York: Churchill Livingstone;1989.p.229-332.

Conference Proceedings:

Pacak K, Aguilera G, Sabban, E, Kvetnansky R, editors. Stress: current neuroendocrine and genetic approaches. 8th symposium on Catecholamines and Other Neurotransmitters in Stress: 2003 Jun 28-July 3; SmoleniceCastle(place of conference), Slovakia. New York(place of publication): New York Academy of Sciences(publisher); 2004 Jun. 590 p.

Scientific and Technical Reports:

Page E, Harney JM. Health hazard evaluation report. Cincinnati(OH)(Place of publication: Natonal Institute for Occupational Safety and Health (US)(Publisher); 2001 Feb.24 p(Total number of pages). Report No.: HETA2000-0139-2824

Dissertation &Thesis:**Entire Reference**

Kempner JL. Aching heads,making medicine: gender and legitimacy in headache [dissertation]. [Philadelphia]: University of Pennsylvania; 2004.271p.

Alam M. Study of Heart Rate Variability in Adolescent Athletes[M Phil thesis].[Dhaka]: Bangabandhu Sheikh Mujib Medical University; 2008. 178p.

Part of Dissertation &Thesis:

Mackowski MP. Human factors: aerospace medicine and the origins of manned space flight in

the United States[dissertation].[Tempe(AZ)]: Arizona State University;2002 May.Part 2,Space medicine;p.188-377.

Alam M. Heart Rate Variability in Adolescent Athletes[M Phil thesis].Dhaka(Bangladesh): Bangabandhu Sheikh Mujib Medical University; 2008 July.Appendix(Name of the part) 4(Number of the part),Classification of Physical Activity Intensity(Title of the part). p.7(Location of the part).

Standard journal article on the Internet:

Kaul S, Diamond GA. Good enough: a primer on the analysis and interpretation of noninferiority-trials. Ann Intern Med [Internet]. 2006Jul4 [cited 2007 Jan 4];145(1):662-9.Available from:<http://www.annals.org/cgi/reprint/145/1/62.pdf>

Journal article on the Internet with organization (s) as author:

National Osteoporosis Foundation of South Africa. Use of generic alendronate in treatment of osteroporosis. S Afr Med J [Internet].2006 Aug[cited 2007 Jan 9];9(8):696-7. Available from: <http://blues.sabinet.co.za/WebZ/Authorize?>

Journal article on the Internet with governmental body as author:

Centers for Disease Control and Prevention(US), National center for HIV/AIDS,Hepatitis,STD, and TB Prevention and control of tuberculosis in correctional and detention facilities: recommendations from CDC. Endorsed by the Advisory Council for the elimination of tuberculosis, the national Commission of Correctional Health Care and the American correctional Association. MMWR R Rep[Internet].2006Jul7[cited2007Jan 9];55(RR-9):1-44.Availablefrom:<http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5509a1.htm>

Journal article on the Internet with no author:

Prevention strategies for asthma-secondary prevention.CMAJ[Internet].2005 Sept[cited 2007 Jan 5];173(6Suppl):S25-7.Available from:http://www.cmaj.ca/cgi/content/full/173_6_suppl/S25

Journal article on the Internet without standard volume, issue or article number:

Jacobs JL, Lee MT, Lindberg M, Kamin C. Problem based learning,multimedia paucity of behavioral issue learning.MedEduc Online[Internet].2005 [cited 2005]: [5p].Available from:<http://www.med-ed-online.org/pdf/10000006.pdf>

The Editors reserves the customary right to style and if necessary shortens the material accepted for publication and to determine the priority and time of publication. Editor assumes that work based on honest observations. It is not a task of the editor to investigate scientific fraud paper.

E-mail: atlincoln71@gmail.com

For Subscription: Annual subscription for Institution : Taka 300.00, Individuals : Taka 200.00, Senior faculty : Taka 200.00, Junior faculty : Taka 100.00, Overseas patrons : US\$ 20.00. Send subscription to: The Editor, PMJ E-mail: amsshasan@gmail.com

What type of personal data do we handle?

There are four main categories of personal data stored by our journal platform, our press platform, and our book management system; Website User data, Author data, Reviewer data and Editor data.

The minimum personal data that are stored are:

- full name
- email address
- affiliation (department, and institution)
- country of residence

Optionally, the user can provide:

- salutation
- associated URL
- phone number
- reviewing interests
- mailing address
- ORCID
- Google Scholar
- profile picture

What do we do to keep that data secure?

We regularly backup our databases, and we use reliable cloud service providers (Amazon, Google Cloud, Linode) to ensure they are kept securely. Backups are regularly rotated and the old data is permanently deleted. We have a clear internal data handling policy, restricting access to the data and backups to key employees only. In case of a data breach, we will report the breach to the affected users, and to the press/journal contacts within 72 hours.

How do we use the data?

Personal information is only used to deliver the services provided by the publisher. Personal data is not shared externally except for author names,

affiliations, emails, and links to ORCID and social media accounts (if provided) in published articles and books which are displayed as part of the article/book and shared externally to indexes and databases. If a journal operates under open peer review then the reviewer details are published alongside the reviewer details.

2. When registering as an author and submitting an article or book

2.1 what data we collect

- When registering an account we ask you to log in and provide certain personal information (such as your name and email address), and there is the option to register using an ORCID which will pre-fill the registration form.
- As part of submitting an article for publication, you will need to provide personally identifying information which will be used for the peer review process, and will be published. This can include ‘Affiliation’ ‘Competing’ ‘interests’, ‘Acknowledgements’.

2.2 why we collect the data

- Registering an account allows you to log in, manage your profile, and participate as an author/reviewer/editor. We use cookies and session information to streamline your use of the website (for example in order for you to remain logged-in when you return to a journal). You can block or delete cookies and still be able to use the websites, although if you do you will then need to enter your username and password to login. In order to take advantage of certain features of the websites, you may also choose to provide us with other personal information, such as your ORCID, but your decision to utilize these features and provide such data will always be voluntary.
- Personal data submitted with the article or book is collected to allow follow good publication ethics during the review process, and will form part of the official published record in order for the provenance of the work to be established, and for the work to be correctly attributed.

2.3 what we do (and don't do) with the data

- We do not share your personal information with third parties, other than as part of providing the publishing service.
- As a registered author in the system you may be contacted by the journal editor to submit another article.

- Any books published on the platform are freely available to download from the publisher website in PDF, EPUB and MOBI formats on the publisher's site.
- Any personal data accompanying an article or a book (that will have been added by the submitting author) is published alongside it. The published data includes the names, affiliations and email addresses of all authors.
- Any articles published on the platform are freely available to download from the publisher website in various formats (e.g. PDF, XML).
- Ubiquity Press books and articles are typeset by SiliconChips and Diacritech. This process involves them receiving the book and book associated metadata and contacting the authors to finalise the layout. Ubiquity Press work with these suppliers to ensure that personal data is only used for the purposes of typesetting and proofing.
- For physical purchases of books on the platform Ubiquity Press use print on demand services via Lightning Source who are responsible for printing and distribution via retailers. (For example; Amazon, Book Repository, Waterstones). Lightning Source's privacy policy and details on data handling can be found on their website.

2.4 why we store the data

- We store the account data so that you may choose to become a reviewer and be able to perform those tasks, or to become an author and submit an article and then track progress of that article.
- Published personal data that accompanies an article or a book forms part of the official published record in order for the provenance of the work to be established, and for the work to be correctly attributed.

2.5 what to do if you want to get a copy of your data, or want your data to be removed

- You are able to view, change and remove your data associated with your profile. Should you choose to completely delete your account, please contact us at support@ubiquitypress.com and we will follow up with your request as soon as possible.
- To conform to publication ethics and best practice any personal data that is published alongside an article or book cannot be

removed. If you have a query about a publication to which you are attributed please contact info@ubiquitypress.com

3. When registering as a reviewer

3.1 what data we collect

- To become a reviewer you must first register as a user on the website, and set your preference that you would like to be considered as a reviewer. No new personal data is collected when a registered user elects to become a reviewer.
- When registering an account we ask you to log in and provide certain personal information (such as your name and email address), and there is the option to register using an ORCID which will pre-fill the registration form.
- Reviewers can also be registered by editors who invite them to review a specific article. This requires the editor to provide the reviewer's First Name, Last Name, and Email address. Normally this will be done as part of the process of inviting you to review the article or book.
- On submitting a review, the reviewer includes a competing interest statement, they may answer questions about the quality of the article, and they will submit their recommendation.

3.2 why we collect the data

- The data entered is used to invite the reviewer to peer review the article or book, and to contact the reviewer during and the review process.
- If you submit a review then the details of your review, including your recommendation, your responses to any review form, your free-form responses, your competing interests statement, and any cover letter are recorded.

3.3 what we do (and don't do) with the data

- This data is not shared publicly and is only accessible by the Editor and system administrators of that journal or press.
- The data will only be used in connection with that journal or press.
- Data that is retained post final decision is kept to conform to publication ethics and best practice, to provide evidence of peer review, and to resolve any disputes relating to the peer review of the article or book.

- For journals or presses that publish the peer reviews, you will be asked to give consent to your review being published, and a subset of the data you have submitted will become part of the published record.

3.4 what to do if you want to get a copy of your data, or want your data to be removed

- If you would no longer like to be registered as a reviewer you can edit your profile and tick the box 'stop being a reviewer'. This will remove you from the reviewer database, however any existing reviews you may have carried out will remain.
- If you have been contacted by an editor to peer review an article this means that you have been registered in the system. If you would not like to be contacted for peer review you can reply to the email requesting that your data be deleted.

4. When being registered as a co-author

4.1 what data we collect

- Co-author data is entered by the submitting author. The submitting author will already have a user account. According to standard publishing practice, the submitting author is responsible for obtaining the consent of their co-authors to be included (including having their personal data included) in the article/book being submitted to the journal/press.
- The requested personal data for co-authors are at the bare minimum; first name, last name, institution, country, email address. This can

also include; ORCID ID, Title, Middle Name, Biographical Statement, Department, Twitter Handle, LinkedIn Profile Name or ImpactStory ID.

4.2 why we collect the data

- Assuming that it is accepted for publication, this data forms part of the official published record in order for the provenance of the work to be established, and for the work to be correctly attributed.
- Author names, affiliations and emails are required for publication and will become part of the permanent cited record.

4.3 what we do (and don't do) with the data

- The co-author's personal data is stored in the author database. This personal data is only used in relation to the publication of the associated article.
- Any co-author data collected is added to the author database and is only used in association with the article the user is co-author on.

Notification about change of ownership or of control of data

We may choose to buy or sell assets. In the case that control of data changes to or from Ubiquity Press and a third party or in the case of change of ownership of Ubiquity Press or of part of the business where the control of personal data is transferred, we will do our best to inform all affected users and present the options.

Artificial Intelligence and Medical Science: Present Perspective

Ahmmad Taous

Introduction

Artificial Intelligence (AI) is a branch of computer science that has extensive and diversified scope of work, concerned with building smart machines capable of performing tasks that typically require human intelligence. Some applications of AI include automated interfaces for visual perception, speech recognition, decision-making and translation between languages. It is an interdisciplinary science.

Extended healthcare services nowadays contains huge amount of data and it rises! So AI becomes more relevant in healthcare sector. Several types of AI are already being employed by payers and providers of healthcare. The key categories of applications are patient engagements and adherence, diagnosis, treatment recommendations, drug discovery as well as administrative activities.

It is widely accepted that the term AI was first thought in 1956 when American computer scientist John McCarthy *et al.* organized the Dartmouth Conference. It came forward when Turing test proposed by Alan Turing as a measure of machine intelligence and a chess-playing program written by Dietrich Prinz.

Artificially intelligent systems in healthcare have the following typical pattern. Such a system starts with a large amount of data; on these data machine-learning algorithms are employed to get information which is to generate a more data to solve a problem in the healthcare system. It also include matching patient symptoms to appropriate physician for their diagnosis, prognosis of disease, drug discovery, translate languages, organize and interpret different images and files.

Discussion

AI systems to apply to drug discovery and doctors personal treatment planning were contents of huge amount of research! Companies, such as Verge Genomics, focus on the application of machine-learning algorithms to analyze human genomic data

and identify drugs to combat neurological diseases, such as Parkinson's, Alzheimer's, amyotrophic lateral sclerosis (ALS) in a cost-effective way.

Many companies such as Bot MD have built systems that can help patients with clinical issues regarding:

- Instantly finding which physicians are on call and scheduling the next available appointment; the AI system can also search multiple scheduling systems across different hospitals
- Answering prescription related questions, like drug availability and cost-effective alternative drugs
- Assisting doctors search hospital protocol, list of available clinical tools and available drugs all through the use of a mobile application, thus improving workflow in the hospital.

The most recent application of AI in global healthcare is the prediction of emerging hotspots using contact tracing and flight traveler data to fight off the novel Coronavirus (COVID-19) pandemic.

Artificially intelligent systems is used for diagnosing some specific disease of patients like dematological conditions which can be diagnosed by physicians visual efficiency, medical data of radiological films are processed and analyzed by artificially intelligent machine system. In ophthalmological fields of research, much work has been done. Researchers at Google developed and trained a system using 128,175 retinal fundus images to classify them into diabetic retinopathy and macular edema for adults with diabetes. Here the AI model shows several advantages such as:

1. Automated grading of diabetic retinopathy leading to increased efficiency in diagnosing many patients in shorter time;
2. AI can give second opinion to the Ophthalmologist as assistant to him!

3. Early detection of diabetic retinopathy due to capability of the model to study images at the granular level which is rarely possible for a human Ophthalmologist!

There is virtually no area in medicine and care delivery that is not already being touched by artificial intelligence!

Contact tracing is a disease control measure used by government authorities to limit spread of a disease. Contact tracing works by contacting and informing individuals that have been exposed to a person who has contacted the disease. Then they are advised to be quarantined or isolated to prevent further spread of the disease. As reported by Apple Newsroom, tech giants like Google and Apple have joined forces to create a contact tracing platform that will use AI systems through the use of application programming interfaces. That interface is programmed in the smartphones to reach the users. Location services of Smartphones are then allowed in the vicinity of the infected person which ultimately reported to government health system.

BlueDot, a Canadian company works in health sector. They first published scientific article on AI and ML that can accurately predicted worldwide spread of the virus. They also used other techniques like natural language processing (NLP). About 100,000 articles from over 65 countries analyzed searching travel information and flight paths, an area's climate, temperature and even local livestock to help predict future outbreaks.

Despite the limitations, AI looks well positioned to revolutionize the healthcare industry. AI systems can help free up the time for busy doctors with follow-up care and availability of prescription drug alternatives. AI also has the capability of remotely diagnosing patients, thus extending medical services to remote areas, beyond the major urban centers of the world. The future of AI in healthcare is bright and promising, and yet much remains to be done.

There are many negative effects of modern technology on mental health. However, researchers at the University of Southern California (USC) in collaboration with Defense Advanced Research Projects Agency and the U.S. Army found that people suffering from post-traumatic stress and other forms of mental anguish are more open to discussing their concerns with virtual humans than actual humans

for fear of judgment. This research has promising results for the role of virtual assistants resulting in the collection of honest answers from patients that could help doctors diagnose and treat their patients more appropriately and with better information.

Most global pharmaceutical companies have invested their time and money on using AI for drug development of major diseases, such as cancer or cardiovascular disease. Apart from these, AI is incorporated into some sleep medicine software solution. This network approach detect sleep pattern more accurately and predict sleep apnoea markers and can score the sleep apnoea indicators. The FDA now incentivizes companies to develop new treatments for these diseases on priority basis. It should be included in medical curriculum because the cost of a wrong decision can be fatal. The data used to learn from and the optimization strategy used has a deep impact on the applicability of the AI system to solve a particular problem.

In the next decades, AI will integrate more with electronic and digital medical records in the analysis of outcome and risk or benefit of any treatment modalities. While the evolving use of artificial intelligence will continue to play a crucial role in patient outcomes, of course the patients treatment plan will always be in human hands

Conclusion

The ongoing development in AI and ML has significantly improved treatment, medication, screening, prediction, forecasting, contact tracing, and drug/vaccine development process for the Covid-19 pandemic and reduce the human intervention in medical practice. Success of AI during this pandemic encouraged the medical experts to use its skill to handle human disease, its diagnosis, risk detection and drug discovery. But it is not widely used by the clinicians, so adoption of AI in clinical practice needs further development.

Pabna Medical College Journal 2023;2(1): 1-2

Professor Dr. Ahmmad Taous

Professor of ENT, Pabna Medical College, Bangladesh
Executive Editor, Pabna Medical College Journal
Editor, Bangladesh Journal of Otorhinolaryngology, Bangladesh
ORCID ID: 0000-00023020-0236
E-mail: atlincoln74@gmail.com

Percutaneous Dilatational tracheostomy: an Observational Study at Birat Medical College and Teaching Hospital, Kathmandu, Nepal

Shah RK¹, Ullah MS², Koirala B³, Rajbansi LK⁴, Hasan DM⁵, Arjyal B⁶

Article info

Received : 21.03.2023
Accepted : 30.04.2023
No. of Tables : 0
No. of Figure : 0
No. of References : 16

Cite the Article:

Shah RK, Ullah MS, Koirala B, Rajbansi LK, Hasan DM, Arjyal B. Percutaneous Dilatational tracheostomy: An observational Study at Birat Medical College and Teaching Hospital. Pabna Med. Cal. J. 2023;2(1): page 3-6.

Keywords: Bleeding, percutaneous dilatation tracheostomy, premature extubation, prolonged intubation

Abstract:

Background: Tracheostomy is the most commonly performed procedure worldwide. It is mainly done in the critically ill patients requiring long term ventilation, facial trauma, laryngeal fractures, laryngeal malignancy, etc. Converting from endotracheal intubation to tracheostomy has many benefits in terms of maintaining the laryngeal function, feeding, safety and it is more comfortable than endotracheal intubation. Tracheostomized patients requires less analgesics and sedatives. It also helps in easier and early weaning from mechanical ventilator, possibility of speech, initiation of oral intake of medication. This all contributes to shorter ICU stay and hospital stay which becomes more cost effective.

Methods: This is a prospective observational study that was conducted in the 22 bedded neuro intensive care and medical intensive care unit of Birat Medical College and Teaching Hospital. The data were collected from 1st Aug 2021 to 28th Feb 2022. All the patients needing tracheostomy for prolonged intubation, protection of airway and to maintain tracheobronchial toileting was taking under study.

Results: Our study was a prospective observational regarding percutaneous dilatational tracheostomy. A total of 71 patients were included in the study. Out of which 39(54.9%) were male and 32(45.1%) were female with the ratio of (1.2:1). The age of the patients ranged from 18-82 years with the mean age of 53.25. The most common indication of percutaneous dilatational tracheostomy was prolonged intubation and that accounted for 29(40.8%) of the patients which was followed by airway protection 27(38%) patients and to maintain the pulmonary hygiene was seen in 15 (21.1%) patients.

Conclusion: Tracheostomy is among the most frequently performed procedures in critically ill patients. Percutaneous dilatational tracheostomy is a safe and feasible procedure performed at bedside with minimal invasive technique in ICU.

Introduction

Tracheostomy is the most commonly performed procedure worldwide.¹ It is mainly done in the critically ill patients requiring long term ventilation, facial trauma, laryngeal fractures, laryngeal

malignancy, etc. Converting from endotracheal intubation to tracheostomy has many benefits in terms of maintaining the laryngeal function, feeding, safety and it is more comfortable than endotracheal intubation. Tracheostomized patients requires less

1. Dr. Rajeev Kumar Shah, Assistant Professor, Department of Otorhinolaryngology, Birat Medical College and Teaching Hospital, Kathmandu, Nepal. ORCID id: <https://orcid.org/0000-0003-2398-157X>
2. Dr. Md. Safi Ullah, Associate Professor (ENT & HNS), Malek Ukil Medical College, Noakhali, Bangladesh.
3. Dr. Bipin Koirala, Assistant Professor, Department of Otorhinolaryngology, Birat Medical College and Teaching Hospital, Nepal
4. Dr. Lalit Kumar Rajbansi, Associate professor, Department of Anaesthesiology and Critical Care, Birat Medical College and Teaching Hospital, Nepal
5. Dr. Dewan Mahmud Hasan, Consultant (ENT & HNS), Impulse Hospital, Dhaka, Bangladesh.
6. Dr. Batsalya Arjyal, Assistant professor,, Department of Anaesthesiology and Critical Care, Birat Medical College and Teaching Hospital, Nepal

Correspondence: Dr. Rajeev Kumar Shah, Assistant Professor, Department of Otorhinolaryngology, Birat Medical College and Teaching Hospital, Nepal. Email- shahrajeev2001@gmail.com, Orchid id: <https://orcid.org/0000-0003-2398-157X>

analgesics and sedatives. It also helps in easier and early weaning from mechanical ventilator, possibility of speech, initiation of oral intake of medication. This all contributes to shorter ICU stay and hospital stay which becomes more cost effective.²⁻⁴The other advantage of percutaneous tracheostomy is it can be done in bedside and there is no need of transferring unstable and critically ill patients to the operating room. This method becomes cost effective and time effective in terms of use of manpower and equipment required.⁵

Shelden and his colleagues first reported percutaneous tracheostomy in 1955.⁶ Several tracheostomy techniques have been described as percutaneous.^{6,7} Several methods of percutaneous techniques came into practice. But the most popular technique which is being practiced was described by Ciaglar⁸ as it performed rapidly and safely at the bedside. In the past open surgical tracheostomy was the only method and is one of the oldest known procedures known to mankind. Only after 1985, percutaneous dilatation tracheostomy was popularized and it became an alternative to traditional open surgical tracheostomy.⁹

The aim of this study is to record the learning curve, operative time, safety, complication and different hemodynamic parameters of the body during the percutaneous tracheostomy.

Methods:

This is a prospective observational study that was conducted in the 22 bedded Neuro Intensive Care and Medical Intensive Care Unit of Birat Medical College and Teaching Hospital, Kathmandu, Nepal. The cases were collected from 1st Aug 2021 to 28th Feb 2022. The written and informed consent for the inclusion in the study was taken from close patient's close relatives. Approval for the study was obtained from Institutional Review Committee.

The inclusion criteria for tracheostomy were patient requiring tracheostomy while the patients having any form of previous radiation of the neck, infection at the site of tracheostomy, age less than 18 years, bleeding diathesis, neck mass, history of neck surgery, patient with cervical injury and emergency tracheostomy were excluded from the study group.

Smith medical portex kit was used for the procedure. The procedure was done under the supervision of an anesthesiologist. Adequate sedation with midazolam

or propofol and adequate analgesia with intravenous fentanyl were given to achieve anesthesia. Routine monitoring including ECG, heart rate, blood pressure and oxygen saturation were done.

The patient was positioned in supine position with head extension which was maintained by sand bag in the back of patient. Before the procedure the tube was pulled back such that the tip of the endotracheal tube was located just at the level below the vocal cord with Murphey's eye below the cord. The tube was then re fixed with tape.

Appropriate size percutaneous dilator (no. 7,7.5,8) was selected for the patient. The site for percutaneous puncture was marked two finger above the sternal notch and was painted and draped. Injection 2% lignocaine with adrenaline (1:80000) was injected at the marked site. A horizontal incision up to the subcutaneous fascia was given and after that artery forcep was used to split underlying tissue. A fourteen-gauge needle was used to locate the trachea and was confirmed by sudden loss of resistance and gush of air in the syringe. Once the needle sheath is inside the trachea, the guide wire was inserted till the white marking on the guidewire followed by A fourteen-gauge needle was used to locate the trachea insertion of the short dilator. The short dilator was removed leaving the guidewire in situ and the long guiding catheter was passed over the guidewire into the trachea in the direction of the arrow marked on the catheter. A well lubricated 'single stage dilator' passed over the guiding catheter until it reached the safety stop. Finally, the dilator was removed and well lubricated tracheostomy tube loaded on its introducer over the guiding catheter through the stoma with a slight twisting motion. The tracheal tube position was then finally confirmed by capnography. After that the tube was connected to mechanical ventilator.

Results:

Our study was a prospective observational regarding percutaneous dilatational tracheostomy. A total of 71 patients was included in the study. Out of which 39(54.9%) were male and 32(45.1%) were female with the ratio of (1.2:1). The age of the patients ranged from 18-82 years with the mean age of 53.25.

The most common indication of percutaneous dilatational tracheostomy was prolonged intubation and that accounted for 29(40.8%) of the patients which was followed by lower airway protection

27(38%) patients and to maintain the tracheo-bronchial toileting was seen in 15 (21.1%) patients.

The intraoperative and post operative complication was noted. Bleeding was the most common complication seen and it accounted for 9(12.7%) patients. This was managed during the interoperative period by applying pressure and suture. Seven patients were managed by pressure while two patients needed suturing. Five patients had stromal site infection. This was because of the excessive secretion and poor tracheostomy care. Stomal site infection was managed with daily dressing and applying dry Neosporin powder. Premature extubation was seen in four patients which was managed by reinsertion of portex tracheostomy tube.

Tracheostomy in faculty passage was seen in 3(4.2%) cases. This complication was noted immediately through capnograph. Transient hypotension was seen in 2 (2.8%) patients who was managed accordingly.

Tubal blockage was seen in 2 (2.8%) cases. This was managed by immediate removal of the tracheostomy and reinsertion of new tracheostomy tube in situ. Two patients had pneumothorax which was managed by insertion of chest tube by surgeons. Similarly, two patients showed tracheoesophageal fistula which was managed conservatively by inserting feeding tube and the patients was kept nil per orally for 7 days. While 42(59.2%) didn't have any kind of complication during the procedure and throughout their hospital stay.

The operating time for percutaneous dilatational tracheostomy was noted. The minimum time noted was 5 minutes while the maximum time noted was 25 minutes. The mean operating time was 11.87. The heart rate ranged from 78-122 beats per minutes with the mean of 93.97. The mean arterial pressure was between 61-110 mm of Hg with the mean of 84.34 during the procedure.

Discussion:

The purpose of our study was to find out the efficacy, operating time, complication and different hemodynamic parameters of the body during the percutaneous dilatational tracheostomy. The main advantage of percutaneous dilatational tracheostomy is the ability to perform it at the bedside in the ICU, thus avoiding a potentially hazardous transfer of critically ill patients to the operating room.

PDT has been shown to reduce significantly the cost of tracheostomy compared with surgical tracheostomy. The incision for percutaneous tracheostomy is relatively smaller compared to open surgical tracheostomy. The portex tracheostomy tube is fitted tightly against the stroma so that there is less tissue dissection and less damage to the tissue in PDT procedure. The advantages being shorter hospital stay and long term follow up studies who underwent PDT in terms of complication and aesthetically more favorable scar.¹⁰⁻¹²

A study conducted by Kearney et al¹³ found out the perioperative complication was seen in 6% of the cases and premature extubation was the most common complication. Early postoperative complication in form of bleeding was seen in 5% of the cases. While in our study bleeding was the most common complication which was seen in 9(12.7%) of the patients. Early postoperative complication in form of bleeding was seen in 5% of the cases. The mortality rate was 0.6%.in the study conducted by Kearney et al.

In the study conducted by Toursarkissian et al¹⁴ the mean procedural time was 15 ± 9 minutes (range 5-60 minutes) while in our study the operating time for the percutaneous dilatational tracheostomy range from 5-25 minutes with the mean of 11.87.

Percutaneous dilatational procedure is a closed procedure and has a unique risk of paratracheal insertion of tracheostomy tube insertion and posterolateral wall laceration. This complication has to be managed surgically with good outcome.¹⁵

There are certain situation in which open surgical tracheostomy is preferred over PDT like inexperienced surgeon, emergency placement of tracheostomy tube, in the condition where the anatomical landmarks are not clearly palpable like short neck, obese, enlarged thyroid gland, nonpalpable cricoid cartilage and gross shifting of the trachea.

Use of adjuncts like preoperative ultrasonography helps in identifying aberrant blood vessels thus preventing damage to the vessels and avoiding risk of bleeding. Based upon the preoperative ultrasound finding the Keling Et al¹⁶ changed the site of tracheal puncture in 24% of the cases. The use of bronchoscopy during the procedure gives additional benefit of confirmation of needle placement, dilatation and tube placement.

Conclusion:

Tracheostomy is among the most frequently performed procedures in critically ill patients. Percutaneous dilatational tracheostomy is a safe and feasible procedure performed at bedside with minimal invasive technique in ICU. It has a very low morbidity if performed by skilled surgeon. The complication following percutaneous dilatational tracheostomy is relatively less. This technique is very cost effective as it can be performed in the bedside thus eliminating the use of operating room and extra man power. It also saves time for transporting the critically ill patients to the operating room. Thus this method is safe, cost effective and can be done in a bed side. It is believed that use of technical adjuncts like use of ultrasound and bronchoscopy will improve patient safety.

References:

1. Frost EA. Tracing the tracheostomy. *Ann Otol Rhinol Laryngol.* 1976 Sep-Oct; 85 (5 Pt.1): 618-24. doi: 10.1177/000348947608500509. PMID: 791052.
2. Hsia, DW, Ghori, UK, Musani, AI. Percutaneous dilatational tracheostomy. *Clin Chest Med.* 2013;34(3):515-526. doi:10.1016/j.ccm.2013.04.002
3. Cheung, NH, Napolitano, LM. Tracheostomy: epidemiology, indications, timing, technique, and outcomes. *Respir Care.* 2014;59(6):895-915; discussion 916-9. doi:10.4187/respcare.02971
4. Durbin, CG. Tracheostomy: why, when, and how? *Respir Care.* 2010;55(8):1056-1068. PMID: 20667153
5. Cobean, R, Beals, M, Moss, C, Bredenberg, CE. Percutaneous dilatational tracheostomy: a safe, cost-effective bedside procedure. *Arch Surg.* 1996;131(3):265-271. doi:10.1001/archsurg.1996.01430150043008
6. Shelden CH, Pudenz RH, Tichy FY. Percutaneous tracheotomy. *J Am Med Assoc.* 1957 Dec 21; 165(16): 2068-70. doi: 10.1001/jama.1957.02980340034009. PMID: 13480860.
7. Toye FJ, Weinstein JD: A percutaneous tracheostomy device. *Surgery* 1969, 65:384-389. doi: 10.1186/cc340. PMID: 11094476
8. P. Ciaglia, Graniero KD. Percutaneous Dilatational Tracheostomy: Results and Long-term Follow-up. *Chest* 101, Issue 2, 1992; 101:464-467. doi:10.1378/chest.101.2.464.
9. Freeman, BD, Isabella, K, Lin, N, Buchman, TG. A meta-analysis of prospective trials comparing percutaneous and surgical tracheostomy in critically ill patients. *Chest.* 2000;118(5):1412-1418. doi: 10.1378/chest.118.5.1412. PMID: 11083694.
10. Stoekli SJ, Breitbach T, Schmid S: A clinical and histologic comparison of percutaneous dilatational versus conventional surgical tracheostomy. *Laryngoscope* 1997, 107: 1643-1646. 10.1097/00005537-199712000-00012
11. Walz MK, Peitgen K, Thurauf N, Trost HA, Wolfhard U, Sander A, Ahmadi C, Eigler FW: Percutaneous dilatational tracheostomy: early results and long-term outcome of 326 critically ill patients. *Intensive Care Med* 1998, 24: 685-690. 10.1007/s001340050645
12. Gysin C, Dulguerov P, Guyot JP, Perneger TV, Abajo B, Chevrolet JC: Percutaneous versus surgical tracheostomy. *Ann Surg* 1999, 230: 708-714. 10.1097/00000658-199911000-00014
13. Kearney PA, Griffen MM, Ochoa JB, Boulanger BR, Tseui BJ, Mentzer RM Jr: A single-center 8-year experience with percutaneous dilatational tracheostomy. *Ann Surg* 2000, 231: 701-709. 10.1097/00000658-200005000-00010
14. Boulos Toursarkissian, Thomas N. Zweng, Paul A. Kearney, Walter E. Pofahl, Steven B. Johnson, Donald E. Barker. Percutaneous dilatational tracheostomy: Report of 141 cases. *The Annals of Thoracic Surgery.* 1994;57(4):862-867. doi.org/10.1016/0003-4975(94)90191-0.
15. Lin JC, Maley RH Jr, Landreneau RJ: Extensive posterior-lateral tracheal laceration complicating percutaneous dilatational tracheostomy. *Ann Thorac Surg* 2000, 70: 1194-1196. 10.1016/S0003-4975(00)01504-6
16. Kollig E, Heydenreich U, Roetman B, Hopf F, Muhr G: Ultrasound and bronchoscopic controlled percutaneous tracheostomy on trauma ICU. *Injury* 2000, 31: 663-668. 10.1016/S0020-1383(00)00094-2

The "Lelli's Test": a Screening Clinical Test for the Diagnosis of Anterior Cruciate Ligament Injury

*Ali MM¹, Naharin NE², Hossain SME³, Manjur M⁴, Islam MN⁵, Masur MAA⁶, Khatun MF⁷

Article info

Received : 03.03.2023
Accepted : 27.04.0023
No. of Tables : 03
No. of Figure : 01
No. of References : 13

Cite the Article:

Ali MM, Naharin NE, Hossain SME, Manjur M, Islam MN, Masur MAA, Khatun MF. The "Lelli's Test": a Screening Clinical Test for the Diagnosis of Anterior Cruciate Ligament Injury. Pabna Med. Cal. J. 2023;2(1): page 7-10.

Keywords: Anterior cruciate ligament, Anterior drawer test, Lachman test, Pivot shift test, Lelli's test

Abstract

Background: The anterior cruciate ligament (ACL) is a major knee stabilizing factor during both of the knee's rotational and anteroposterior stability. ACL damage is a major injury to the knee joint. For identification of ACL injury routinely we practice Anterior drawer test, Lachman's test and Pivot shift test. In this study we evaluate the accuracy of Lelli's test for detecting ACL tear and compare it with MRI findings.

Methods: This study included 100 patients attending outdoor patient department at Department of orthopedics, Combined Military Hospital (CMH) Bogura from February 2022 to February 2023. aged between 20-60 years with knee symptoms of giving way/locking/pain. Clinical history and detail clinical assessment of knee for instability including Lelli's test were performed by standard technique and recorded in the Performa. All patients send for MRI of knee and result were compared with Lelli's test to assess the reliability of clinical test as screening test for ACL injury.

Results: In this study total numbers of cases were 100 with mean age was 35 ± 09 years and most of the patients were male 98(98.00%). Maximum injuries occurred on the right side 80 (80.00%). The most common cause of ACL injury was sports activities 85(85.00%) and chief complaints of giving way 82(82.00%). In this study Lelli's test had sensitivity 94.18 % (CI 70.93-94.57) and Specificity 93.33 % (CI 79.85-97.17)

Conclusion: Lelli's test is a simple with high sensitivity and specificity which can be used as a screening test for evaluation of ACL function in both acute and chronic knee injury.

Introduction

The anterior cruciate ligament (ACL) is a major biomechanical component of knee stabilizing factor during both of the knee's rotational and anteroposterior stability.¹ Anterior cruciate ligament (ACL) damage is a major injury to the knee joint,² with an annual incidence of 68.6 per 100,000 person-years and.³ In the United States nearly 200,000 injuries of ACL occur per year, with an estimated 100,000 reconstructions annually.⁴

ACL injuries can be identified by a detail history of knee trauma followed by a typical physical examination finding of anterior sliding of the tibia over the femur.⁵

Anterior drawer test, Lachman's test, pivot shift test, and the Lelli's test are the physical exam tests for detecting ACL deficiency. Compared to the other physical examination maneuvers, the Lelli's test is the simplest and easiest to perform regardless of who performs the test.^{6,7} Lever Sign test would be

1. Dr. Md. Mahbub Ali, Classified Specialist, Department of Orthopaedic Surgery, Combined Military Hospital, Bogura.
 2. Dr. Noor-E-Naharin, Associate Professor, Department of Obstetrics and Gynaecology, Army Medical College, Bogura.
 3. Dr. SM Eqbal Hossain, Classified Specialist, Department of Orthopaedic Surgery, Combined Military Hospital, Barishal.
 4. Dr. Mushfique Manjur, Assistant professor, Department of Orthopaedic Surgery, Monno medical college & Hospital.
 5. Dr. Md Nazrul Islam, Junior consultant, Department of Orthopaedic Surgery, Kalai Upazilla Health Complex, Kalai, Joypurhat.
 6. Dr. Mohammad Akshad Al Masur, Senior consultant, Department of Orthopaedic Surgery, 250 Bedded General Hospital, Pabna.
 7. Dr. Mst. Fatema Khatun, Medical Officer, Department of Obstetrics and Gynaecology, 250 Bedded General Hospital, Pabna.
- Correspondence:** Dr. Md. Mahbub Ali, Classified Specialist, Department of Orthopaedic Surgery, Combined Military Hospital, Bogura. E-mail- mahbub101164@gmail.com, Phone- +880 1714084012

diagnostic for both partial and complete tears of the ACL regardless of the elapsed time from injury.⁸

For diagnosis of ACL injury arthroscopic visualization is the gold standard technique and magnetic resonance imaging (MRI) is a valid and noninvasive diagnostic method, with a specificity and sensitivity of 94–98 %.⁸

Since the Lelli's test is simple, it can be performed with high accuracy by all medical personnel, even if done by non-orthopedic providers. Our aim in this study is to measure the accuracy of the test to performed as a screening test for ACL injury.¹

Methods

This survey was carried out in CMH Bogura, a tertiary hospital of Bangladesh from February 2022 to February 2023. Total 100 patients included in this study who were attending orthopaedic outpatient department, aged between 20-60 years with knee symptoms of giving way or locking or pain following sports or non-sports injury. Clinical history and detail clinical assessment of knee for instability including Anterior drawer test, Lachman test, Pivot shift test, Lelli's test were performed by standard technique and recorded in the Performa. Lelli's test was carried out with patient in supine position, knee in full extension and heel touching the bed, clenched fist is placed

beneath the proximal calf just distal to tibial tuberosity then a gentle posterior directed thrust is applied over quadriceps tendon and we look for heel liftoff from the bed. The test is positive if there is no heel liftoff from the bed in springy manner compared to normal side. All patients send for MRL of knee and result were compared with Lelli's test finding

Ethical permission had been taken from institute keeping compliance with Helsinki Declaration for medical research involving human subject 1964. The study subjects were informed verbally about the study design, the purpose of the study and right for withdrawing themselves from the project at any time, for any reason, what so ever. Persons who gave informed written consent to participate voluntarily in the study were included in the study sample. The SPSS version 20.0 was used for data analysis.

Results

In this study total numbers of cases were 100. 35 ± 09 years was the mean age of the patients. Among them, 02% ($n=2$) were female and 98% ($n=98$) were male. Maximum injuries occurred on the right side 80 % ($n=80$). The most common cause of ACL injury was sports activities 85(85.00%) and chief complaints of patients were giving way 82(82.00%), pain 43(43.00%), locking 28(28.00%). Demographic and clinical features of the patients are given in Table I, II.




Figure - 1: (A) A photograph of the lever test setup, with the examiner's hand positioned for the test. (B) A photograph of a negative lever test, with the femur being translated posteriorly using the right hand (downward red arrow) and levering of the tibia resulting in lift-off of the heel (upward red arrow). The examiner's fist is just distal to the tibia tubercle. (C) A photograph of a positive lever test, with the femur being translated posteriorly using the left hand (downward red arrow) and translation of the proximal tibia anteriorly (white upward arrow) with no heel lift-off¹.

Table I: Demographics features of the survey respondents (n=100)

Demographic Feature			
Age		Mean \pm SD	35 \pm 09 yrs
Gender	Female	N(%)	2(02.00%)
	Male	N(%)	98(98.00%)
Lesion	Right	N(%)	80(80.00%)
	Lt	N(%)	20(20.00%)
Cause	Sports injury	N(%)	85(85.00%)
	Fall on slippery ground	N(%)	02(02.00%)
	Fall from height	N(%)	07(07.00%)
	Trauma	N(%)	06(06.00%)

Table II: Clinical Features of the survey respondents (n=100)

Clinical Features		
Only giving way	N (%)	37(37.00%)
Only Pain	N (%)	05(05.00%)
Only Locking	N(%)	10(10.00%)
Both giving way and pain	N(%)	30(30.00%)
Both giving way and locking	N(%)	10(10.00%)
Both locking and pain	N(%)	03(03.00%)
All symptoms including giving way, pain and locking	N(%)	05(05.00%)

Clinical diagnosis of ACL injury

Comparison of the MRI findings of ACL Tear and clinical Lelli's test findings yielded the following results shown in table-3.

Table III: Lelli's test and ACL tear

Lelli's test	MRI findings		Total	Sensitivity (CI)	Specificity (CI)
	ACL tear present	ACL tear absent			
Positive	80	1	81	94.18(70.93-94.57)	93.33(79.85-97.17)
Negative	5	14	19		
Total	85	15	100		

Lelli test had sensitivity 94.18 % (CI 70.93-94.57) and Specificity 93.33 % (CI 79.85-97.17)

Discussion

In most of the patients excess knee laxity and the joint becomes unstable is the most detectable signs after ACL rupture.⁹ Anterior drawer, the Lachman, and the pivot shift test are the three commonly applied tests to determine ACL injuries.¹⁰

In a meta-analysis of Twenty-eight studies that assessed the accuracy of clinical tests for diagnosing ACL ruptures by Benjaminse et al² with pooled sensitivity of Anterior drawer test, Lachman test, Pivot shift test 92% (CI 88-95) , 85% (CI 83-87), 24% (CI 21-27) respectively and pooled specificity 91% (

CI 87-94) ,94% (95% CI, 92-95), 98% (CI 96-99) respectively.

In this study we found Lelli test had sensitivity 94.18 % (CI 70.93-94.57) and Specificity 93.33 % (CI 79.85-97.17), which is comparable to three commonly applied tests in clinical practice to determine ACL injuries. There are limitations to the commonly applied test like for Lachman test; examiners who have small hands may face difficulties on patients with a large thigh girth². For pivot shift test patient with a chronic ACL-deficient knee is familiar with unpleasant phenomenon of pivoting and will show protective muscle action.¹¹ The accuracy of a complex test maneuver such as the pivot shift test may increase with experience thus the pivot shift test has very high specificity and low sensitivity. For Anterior drawer test the hemarthrosis and reactive synovitis may preclude knee flexion to 90°, hindering the proper performance of the test, protective muscle action of the hamstrings secondary to joint pain provides a vector force opposite to the anterior translation of the tibia and the posterior horn of the medial meniscus becomes buttressed against the posterior most margin of the medial femoral condyle and may preclude anterior translation of the tibia¹². We found Lelli's test to be least affected by those phenomenon, relatively easy to carryout in both acute and chronic knee injury situation with comparable sensitivity and specificity to those routinely applied clinical test for ACL tear. The most significant finding of this study was the perfect agreement between the outcome of the Lever Sign test and the MRI findings, regardless of whether the ACL was partially or completely ruptured and regardless of the time elapsed from the injury. MR has been shown to have a specificity and sensitivity of 94–98 % in detecting ACL injury^{2,12,13}.

Conclusion

Lelli's test is a simple with high sensitivity and specificity in comparison to those routinely practiced clinical test for ACL injury. As such it can be used as a screening test for evaluation of ACL function in both acute and chronic knee injury. But for more validity a multi-centric meta-analysis is required.

Conflict of interests: None declared.

References

1. Dawod MD, Alswerki MN, Darabah AJ, Darabah YJ, Akel AY, Alisi MS. Clinical Reproducibility and Reliability of Lever Sign (Lelli's) Test for Acute ACL

Tear Performed by Medical Students. *Advances in Medical Education and Practice*. 2023 Dec 31;289-95.

2. Benjaminse A, Gokeler A, van der Schans CP. Clinical diagnosis of an anterior cruciate ligament rupture: a meta-analysis. *Journal of orthopaedic & sports physical therapy*. 2006 May;36(5):267-88.
3. Chong AC, Whitetree C, Priddy MC, Zimmerman PR, Haeder PR, Prohaska DJ. Evaluating different clinical diagnosis of anterior cruciate ligament ruptures in providers with different training backgrounds. *The Iowa orthopaedic journal*. 2017;37:71.
4. Jarbo KA, Hartigan DE, Scott KL, Patel KA, Chhabra A. Accuracy of the lever sign test in the diagnosis of anterior cruciate ligament injuries. *Orthopaedic journal of sports medicine*. 2017 Oct 9;5(10):2325967117729809.
5. Parwaiz H, Teo AQ, Servant C. Anterior cruciate ligament injury: A persistently difficult diagnosis. *The Knee*. 2016 Jan 1;23(1):116-20.
6. Gürpýnar T, Polat B, Polat AE, Çarkçý E, Öztürkmen Y. Diagnostic accuracy of lever sign test in acute, chronic, and postreconstructive ACL injuries. *BioMed Research International*. 2019 Jun 9;2019.
7. Lange T, Freiberg A, Dröge P, Lützner J, Schmitt J, Kopkow C. The reliability of physical examination tests for the diagnosis of anterior cruciate ligament rupture—a systematic review. *Manual therapy*. 2015 Jun 1;20(3):402-11.
8. Lelli A, Di Turi RP, Spenciner DB, Dòmini M. The "Lever Sign": a new clinical test for the diagnosis of anterior cruciate ligament rupture. *Knee surgery, sports traumatology, arthroscopy*. 2016 Sep;24:2794-7.
9. Shelbourne KD. The art of the knee examination: where has it gone?. *JBJS*. 2010 Aug 4;92(9):e9.
10. Makhmalbaf H, Moradi A, Ganji S, Omidi-Kashani F. Accuracy of lachman and anterior drawer tests for anterior cruciate ligament injuries. *Archives of bone and joint surgery*. 2013 Dec;1(2):94.
11. SLOCUM DB, JAMES SL, LARSON RL, SINGER KM. Clinical test for anterolateral rotary instability of the knee. *Clinical Orthopaedics and Related Research*®. 1976 Jul 1(118):63-9.
12. Torg JS, Conrad W, Kalen V. Clinical I diagnosis of anterior cruciate ligament instability in the athlete. *The American journal of sports medicine*. 1976 Mar;4(2):84-93.
13. Miyasaka KC. The incidence of knee ligament injuries in the general population. *Am J Knee Surg*. 1991;4:3-8.

Factors Related to Respiratory Distress Syndrome in Neonates

*Akhund M¹, Hamid S², Hossain I³, Alam UK⁴, Noor IN⁵

Article info

Received : 19.02.2023
Accepted : 15.04.0023
No. of Tables : 3
No. of Figure : 0
No. of References : 29

Cite the Article:

Akhund M, Hamid S, Hossain I, Alam UK, Noor IN. Factors Related to Respiratory Distress Syndrome in Neonates. Pabna Med. Col. J. 2023;2(1): 11-17.

Keywords: Respiratory distress syndrome, neonatal mortality, intercostals recession, intercostals recession.

Abstract

Background: Respiratory distress syndrome of neonates (RDS) is one of the most common causes of neonatal mortality.

Objective: The purpose of this study was to investigate socio-demographic, maternal and neonatal factors related to RDS in neonates in context of tertiary level hospital.

Methods: This cross sectional study was conducted in Dhaka Medical College Hospital and Ad-din Women's Medical College and Hospital during 1st January to 31st December 2017 among 186 purposively selected neonates. Data were collected by document review and face to face interview with complete privacy using semi-structured questionnaire and check list.

Results: Regarding socio-demographic factors, Considering the maternal factors, majority of the mothers (61.3%) had taken 4 or more checkup during their antenatal period and 87.1% had institutional delivery while approximately half (52.7%) of the mothers had Caesarean delivery. Among the mothers, 17.2% had history of high blood pressure during pregnancy, 10.8% diabetes mellitus, 14% oligohydramnios and 37.1% mothers had PROM. RDS was more common in preterm neonates or below 37 weeks (85.5%) and maximum were low birth weight or below 2500 gram (90.3%), 52 babies (28%) were born of multiple pregnancy, more than half of the babies (62.4%) had moaning sound (grunting) and most of all (92.5%) had chest recession.

Conclusion: The factors related to RDS are not preventable but are predictable if ANC for all the mothers can be ensured. If the possibility of occurrence of RDS can be assumed considering these factors, necessary precautions for earlier diagnosis and adequate management can be taken which can contribute in reducing neonatal mortality rate.

Keywords: Respiratory distress syndrome, neonatal mortality, intercostals recession, intercostals recession.

Introduction

Infections and complications during the first 28 days of life are responsible for approximately 1.5% of total child deaths. Within this period of life 7% to 50% of neonates are likely to suffer from RDS. Among them 3 to 4 out of 10 requires admission¹. Statistics show that even in USA every year 20,000 to 30,000 neonates suffer from RDS and about 1% pregnancies have got the possibility to have this complication.

RDS is the second most common reason for the respiratory distress in newborns and those who suffer from respiratory distress are 2 to 4 times more prone to death from those who do not².

About 10 -12% of total live born babies are born before completion of their full term gestation³. Globally, in 2010 more than one out of 10 babies was born before completion of 37 weeks of intrauterine life. Again, in the same year around one million babies died due to

1. Dr. Moulee Akhund, Clinical Fellow, ICDDR, Dhaka.
2. Dr. Shaheda Hamid, Associate Professor, Dept. of Maternal and Child Health, National Institute of Preventive & Social Medicine (NIPSOM), Mohakhali, Dhaka-1212.
3. Dr. Irin Hossain, Assistant Professor and Head of the department (OEH).
4. Dr. Ummul Khair Alam, Assistant Professor, Reproductive & Child Health NIPSOM, Mohakhali.
5. Dr. Irfan Nowroze Noor, OSD, DGHS, Doctoral Student, ASEAN Institute for Health Development, Mahidol University, Thailand.

Correspondence: Dr. Moulee Akhund, Clinical Fellow, ICDDR, Dhaka, Phone: 01728419741, E-mail: mouleeakhund@gmail.com

this preterm condition⁴. At present, preterm delivery is the second leading cause of death among under 5 children and the single most important cause of neonatal death⁵. About 50% of babies who are born at 26–28 weeks of gestation are likely to develop RDS while the risk is about only 25% at 30–31 weeks of gestational age. Among the neonates born in 37 weeks of gestation, only 7.8% suffer from RDS⁶.

Birth weight of the baby is considered to be another important concern regarding development of RDS. Different studies found that among the babies born with a weight of 2500 gram or more proportion of developing RDS varies from 9.9%⁶ to 11.5%⁷. Moreover, a previous study shows that proportion of developing RDS is inversely related to the birth weight of the newborns⁸.

Again, according to Nelson Text Book of Paediatrics⁹ not only neonatal factors but also some maternal factors influence in both occurrence and outcome of RDS. The second born of preterm twins⁹, mature neonates with meconium aspiration syndrome¹⁰, delayed crying or perinatal asphyxia (hypoxic ischaemic encephalopathy)¹¹, congenital anomalies¹⁰, IDM (Infant of diabetic mothers)¹², delivery by LUCS, particularly in absence of labour¹³ are commonly documented risk factors in developing RDS among the neonates found in different countries in different studies.

Apnea (cessation of breathing), cyanosis (bluish or blackish discolouration of body), grunting (a moaning sound at expiration), inspiratory stridor, nasal flaring, poor feeding, tachypnea (respiratory rate 60 or more), lower chest wall indrawing (intercostal, subcostal, or suprasternal recession) are considered as the more commonly presented features in a neonate suffering from RDS. A neonate suffering from RDS may present with any one or more than one of the aforesaid clinical manifestations either just after birth or within a very short time after birth, usually with increasing severity with the course of time, reaching the worst peak within the first 48-72 hours of life¹⁴.

Despite all new options like mechanical ventilators and continuous positive airway pressure (CPAP), mortality rates still remain as high about 36%. In a retrospective study carried out in Neonatal Intensive Care Unit (NICU) of Ad-din Hospital from July 2013 to June 2014, it was found that among all indications for mechanical ventilation, RDS contributed to the maximum which was 62% followed by 23% mortality

rate¹⁵. In another retrospective study conducted from December 2014 to November 2015 at the neonatal unit of North East Medical College and Hospital in Sylhet, Bangladesh, it was observed that RDS was responsible for 0.6% of admission into hospital and contributed to 16.7% of the total mortality rate¹⁶.

In a bulletin of the World Health Organization, neonatal deaths presently contribute to more than two-thirds of under-one mortality and for about half of all deaths in children under-five years of age¹⁷. Neonatal mortality rate (NMR) is 28 per 1,000 live births in Bangladesh¹⁸. So, proper measures must be taken to promote the survival of children under five years of age because by signing in SDG, the Government of Bangladesh has declared the vision of decreasing the mortality of children below five years of age by two thirds.

A previous study in Bangladesh¹⁵ stated that although RDS is the most common reason for cardio-respiratory failure in newborns, it also provides with the best prognosis compared to the other reasons if measures can be taken timely. Early intervention can significantly decrease the complications, severity, requirement for oxygen therapy and ventilation, duration of hospital stay¹⁹ and ultimately treatment cost.

Methods:

This cross sectional study was done from 1st January to 31st December 2017, which was approved by the ethical review board of National Institute of Preventive and Social Medicine. Data was collected through face to face interview from purposively chosen 186 mothers of admitted newborn babies diagnosed as RDS into Dhaka Medical College Hospital and Ad-din Women's Medical College and Hospital. Document review was done to collect the data about ANC visits and obstetric history of the mothers as well as the health related events of the neonates since birth. Confidentiality of the respondents was maintained with outmost care and treatment of the neonates in acute condition was ensured prior the data collection.

Statistical Package for Social Science (SPSS) version 20 for windows was used to analyze the data. Associations among qualitative and quantitative variables of various socio-demographic, maternal and neonatal factors were studied by using chi-square test and t-test where appropriate. To assess the independent relationship within the factors, logistic regression analysis was performed. A p value < 0.05 was considered to be significant.

Results:**Table I:** Distribution of the neonates by baseline characteristics and various neonatal factors (n=186)

Gestational Age	Frequency (n)	Percent (%)
24-27 weeks	9	4.8
28-30 weeks	37	19.9
31-32 weeks	37	19.9
33-34 weeks	38	20.4
35-36 weeks	38	20.4
37-39 weeks	27	14.5
Total	186	100.0
Birth weight in gram		
<750	4	2.2
750-1499	61	32.8
1500-2499	103	55.4
2500-3900	18	9.7
Total	186	100.0
Sex		
Male	128	68.8
Female	58	31.2
Cyanosis at Birth		
Present	97	52.2
Absent	74	39.8
Record not Found	15	8.1
Grunting		
Present	116	62.4%
Absent	70	37.6%
Subcostal Recession		
Present	172	92.5%
Absent	14	7.5%
History of delayed cry after birth		
Yes	34	18.3
No	150	80.6
Record not found	2	1.1
Respiratory rate of the baby on admission		
Absent/gasping	5	2.7
<60/minute	55	29.6
>60/minute	126	67.7
Less Foetal Movement		
Yes	34	18.3
No	150	80.6
Don't know	2	1.1
Treatment modality used for giving oxygen		
Room air	32	17.2
Oxygen with nasal prong	11	5.9
Oxygen with face mask	71	38.2
Oxygen with hood	37	19.9
Bubble CPAP	17	9.1
Mechanical ventilation	18	9.7
Severity of RDS		
Less Severe	151	81.2%
More Severe	35	18.8%

Table II: Distribution of the mothers by socio-demographic characteristics and maternal factors (n=186)

	Frequency (n)	Percent (%)
Maternal Age in years		
Below 19	40	21.5
20-24	63	33.9
25-29	45	24.2
30-34	25	13.4
35-39	12	6.5
40-45	1	0.5
Educational qualification of mothers		
Below Primary	42	22.6
Below Secondary	69	37.1
Higher Secondary and Under	45	24.2
Graduate		
Graduate and Post Graduate	30	16.1
Occupation		
Home maker	127	68.3
Teacher	16	8.6
Service Holder	21	11.3
Business	11	5.9
Others	11	5.9
Area of Residence		
Rural	103	55.4%
Urban	83	44.6%
Number of antenatal check up during pregnancy		
No ANC	1	0.5
<4	71	38.2
4 or more	114	61.3
Place of Delivery		
Institution	162	87.1%
Home	24	12.9%
Mode of Delivery		
Spontaneous vaginal	30	16.1
Assisted vaginal	58	31.2
Caesarean	98	52.7
Maternal Factors		
PIH	32	17.2%
GDM	20	10.8%
Oligohydramnios	26	14%
PROM	69	37.1%
Pre-eclampsia	24	12.9%
Placenta Praevia	7	3.8%
Abruptio Placenta	4	2.2%
Multiple Pregnancy	52	28%

Table III: Multivariate Analysis to see the relationship between the socio-demographic variables and number of ANC

Variable	B	P Value	OR	95% C.I. for OR
Rural versus Urban	1.038	0.004	2.823	1.390-5.736
Group of income	-0.185	0.658	0.831	0.367-1.882
Category of occupation	0.034	0.936	1.035	0.453-2.360
Educational Status	1.254	0.003*	3.505	1.543-7.960
Constant	-0.336	0.425	0.715	

*Statistically significant

Discussion

Among the 186 babies, there were 85.5% preterm (below 37 weeks) and 14.5% term neonates (above 37 weeks). So it does not correspond with the traditional saying "RDS only occurs in preterm neonates"¹⁹.

A retrospective case-control study conducted by Liu J et al²⁰ showed that elective LUCS, severe perinatal asphyxia, PROM (Premature Rupture of Membrane), male sex, GDM (Gestational Diabetes Mellitus), LBW (Low Birth Weight), IUGR (Intra Uterine Growth Retardation) and maternal-fetal infection are associated with developing RDS in term newborns. Also, Ghafoor T et al²¹ conducted a prospective study which revealed that 93.61% preterm and 6.38% term neonates developed RDS. The frequency of RDS was, 100% at 26 or less weeks of gestational age, 57.14% at 32 weeks, and 3.70% at 36 weeks of gestational age. Another retrospective cohort study conducted by Mehrabadi et al²² revealed that the rate of RDS per 1000 live births, among live born newborns whose gestational age was <32 weeks and among newborns whose gestational age was 39 weeks or more were 6.4, 113.2 and 1.6 per 1000 live births correspondingly. These results have similarity with the present study that RDS can occur also among term newborns.

More than half (55.4%) weighed between 1500-2499 gram, 61 babies (32.8%) weighed between 750-1499 gram, 18 babies (9.7%) weighed between 2500-3900 gram and only 4 babies (2.2%) weighed <750 gram. In a report prepared by Pramanik A K⁸, the incidence rate of RDS in neonates with birth weight 501-1500g, 501-750g, 751-1000g, 1001-1250g, and 1251-1500g were 42%, 71%, 54%, 36% and 22% correspondingly.

More than two third (68.8%) was boys and 31.2% were girls. This finding corresponds to the result of a study conducted by Ghafoor T et al²¹ which found

that 34.04% were girls and 65.95% were boys. Also, a survey conducted by Luerti M et al²³ revealed that the risk of RDS was more in males than in female neonates.

The more than half of the babies (62.4%) had grunting and the majority (92.5%) had chest recession. A retrospective study conducted by Desai et al²⁴ also supports this finding which revealed that most common clinical manifestations in neonates with RDS are subcostal (xiphoid) recession (98.61%), grunting (93.05%) and intercostal recession (83.33%).

The mean age of the mothers was 24.45 ± 5.627 years and 63 mothers (33.9%) were in the age group of 20-24 years of age. Age specific fertility rate of Bangladesh is maximum (143/1000 women) in this age group¹⁸.

Mothers were almost all educated and among them 37.1% were educated up to secondary level of education. This may be due to the fact that educated mothers are more conscious and more likely to take their babies to the hospital than the mothers educated below the secondary level. More than two third of the mothers (68.3%) were home makers. This reflects the common picture of the females in Bangladesh¹⁸. Most of the above data depict that the sample is similar to national population. Among 186 mothers, maximum proportion i.e. 103 (55.4%) resided in rural area and the rest 83 (44.6%) were inhabitants of urban area. Antenatal care (ANC) from a medically-trained health care provider is essential to evaluate the status of a pregnancy, identify the complications related to the pregnancy and prevent unwanted situations. The present study shows that the majority (61.3%) mothers had taken 4 or more ANC which corresponds with the normal scenario of the women in Bangladesh which states that the majority of women (64 percent) received ANC¹⁸.

Although in Bangladesh, only 38 percent delivery occurs in a health facility, and 62 percent at home¹⁸, this study shows that among the 186 participants, majority (87.1%) of the mothers had institutional delivery whereas only 12.9% had home delivery. This may be due to the fact that those mothers who conduct their delivery at institution have their babies early diagnosed and are conscious enough to bring their child to hospital.

More than half (52.7%) of the mothers had LUCS whereas among the rest 31.2% mothers had assisted vaginal delivery and 16.1% mothers had spontaneous vaginal delivery. Although in Bangladesh, 23 percent of births are found to be delivered by LUCS¹⁸, which implies that six in ten births in a health facility are delivered by C-section. This may be due to the fact that RDS is more prevalent in babies born by LUCS. This finding is supported by the multiple logistic regression statistical method by Bryan H *et al*²⁵ which revealed that the risk of RDS is more with LUCS in absence of labor pain compared to NVD ($p = 0.03$). This result is also similar to a retrospective study conducted by Wu XJ *et al*.²⁶ which revealed that severe RDS can occur in term newborns after elective LUCS. In another study conducted by Goldberg JD *et al*²⁷ among 236 newborns delivered by LUCS, the presence of labour before LUCS did not decrease the risk of RDS significantly, although a trend, not statistically significant, was noted in this direction.

Another hospital-based case-control study conducted by Khalid A. Y²⁸ revealed that the likelihood of LUCS was nearly twice as high among neonates diagnosed with HMD compared to the non-HMD control group. All these information suggest that the chance of RDS is associated with LUCS. Physiology also supports that the vaginal route of delivery is till now known as a better option for enhancement of neonatal cardio-respiratory adaptation²⁸.

This study has also found that out of 186 mothers, 32 (17.2%) mothers had history of PIH, 20 (10.8%) mothers had history of GDM, 26 (14%) mothers had history of oligohydramnios, 69 (37.1%) mothers had history of PROM and 52 babies (28%) were born of multiple pregnancy. This finding is supported by the results of another prospective, descriptive, cross sectional hospital based study, conducted by Swarnkar K and Swarnkar M² which revealed that maternal causative factors like PROM (25.9%), PIH

(16.7%), diabetes mellitus (4.2%) and twin pregnancy (16.7%) were present in cases of RDS. Bowen JR *et al*.²⁹ showed by a case control study that the rate of RDS was more among neonates of mothers with PIH comparison to controls (60% versus 33%; $p = 0.001$).

Besides, the number of ANC visits, education level and wealth status have a positive relationship with the likelihood of delivering in a health facility¹⁸. In Bangladesh half of women who have secondary or higher education and those who are in the higher socio-economic group undergo delivery by LUCS¹⁸.

By logistic regression analysis, the educational status of mothers independently predicted the variation of number of ANC ($p=0.003$, OR=3.505), place of delivery ($p=0.020$, OR=15.925,) and mode of delivery ($p=0.003$, OR=3.505).

To determine the factors influencing the course of RDS, the babies who required mechanical ventilation or bubble CPAP were graded as more severe and the babies who were kept in room air, face mask or oxyhood were graded as less severe RDS. By logistic regression analysis, category of mothers according to income independently predicted the variation of severity of RDS ($p=0.021$, OR=0.379).

Conclusion

This study found that mothers of neonates with RDS share some certain socio-demographic characteristics such as mostly residing in rural area, belonging to lower socio-economic group, educated below secondary level and mostly home maker in occupation. Certain pregnancy associated complications such as diabetes, PIH, oligohydramnios and PROM were common. RDS was more prevalent among male newborns than female ones and neonates born of multiple pregnancy. So as soon as a pregnant mother is diagnosed to have a male foetus or multiple gestation or any of the above mentioned factors she and her family must be prepared for emergency management of the newborn after delivery and transport facilities to higher referral centre must be ensured.

References

1. Saboute M., Kashaki M., Bordbar A., Khalessi N., Farahani Z. The Incidence of Respiratory Distress Syndrome among Preterm Infants Admitted to Neonatal Intensive Care Unit: A Retrospective Study. *Open Journal of Pediatrics*. 2015; 5:285-289.

2. Swarnkar K and Swarnkar M. Neonatal respiratory distress in early neonatal period and its outcome. *International Journal of Biomedical and Advance Research*. 2015;6(9): 643-647.
3. Bhakta KY. Ch 24 A Respiratory distress syndrome. In: John P Cloherty, Ericb C Echeinwald, Ann R Stark. *Manual of neonatal care*. 2010;3: 323-330
4. Blencowe, H., Cousens, S., Oestergaard, M., Chou, D., Moller, A.B., et al. National, regional and worldwide estimates of preterm birth rates in the year 2010 with time trends for selected countries since 1990: a systematic analysis. *The Lancet*, in press; 2012
5. Howson CP, Kinney MV, Lawn JE. *Born Too Soon: The Global Action Report on Preterm Birth*. March of Dimes, PMNCH, Save the Children, WHO. World Health Organization. Geneva 2012. 2012
6. Dani C., Reali M. F., Bertini G., Wiechmann L., Spagnolo A., Tangucci M., Rubaltelli F.F. and the Italian Group of Neonatal Pneumology. Risk factors for the development of respiratory distress syndrome and transient tachypnoea in newborn infants. *European Respiratory Journals Ltd*. 1999.; 14:155-159
7. Ersch J, Roth-Kleiner M, Baeckert P, Bucher HU. Increasing incidence of respiratory distress in neonates. *Acta Paediatr* . 96. 200;96:577-81.
8. Pramanik AK, Rosenkrantz T. *Respiratory Distress Syndrome*. Neonatal Research Network. 2015. Available from: Medscape.
9. Carlo WA and Ambalavanan N. Ch 101.3 Respiratory distress syndrome (Hyaline Membrane Disease). In: *Nelson textbook of Pediatrics*. first South Asia edition. India: Thomson Press India Ltd. 2014;12: 850-858.
10. Clark RH. The epidemiology of respiratory failure in neonates born at an estimated gestational age of 34 weeks or more. *J Perinatol*. 2005; 25:251-7.
11. James LS. Physiology of respiration in newborn infants and in the respiratory distress syndrome. *Pediatrics*. 1959; 12: 1069-101.
12. Robert MF, Neff RK, Hubbell JP, Taeusch HW, Avery ME. Association between maternal diabetes and the respiratory-distress syndrome in the newborn. *N Engl J Med*. 1976; 12:357-60.
13. Hales KA, Morgan MA, Thurnau GR. Influence of labor and route of delivery on the frequency of respiratory morbidity in term neonates. *Int J Gynaecol Obstet*. 1993; 43: 35-40.
14. Morris, I. and Adappa, R. Minimizing the Risk of Respiratory Distress Syndrome. *Paediatrics and Child Health*. 2012. 14:513-517.
15. Mannan MA, Jahan N, Iqbal S, Ferdous N, Dey S, Farhana T and Nazma N. Short Term Outcome of Preterm Neonates Required Mechanical Ventilation. *Chaattagram Maa-O- Shishu Hospital Medical College Journal*. 2016;15(2):9-13.
16. Hamid F, Quaium S M M, Rahman A, Ahmad ATR, Khan S, Hussain T, Zahra F, Islam TMT. Audit of Neonatal Morbidity and Mortality in A Tertiary Care Hospital, Sylhet. *Chaattagram Maa-O-Shishu Hospital Medical College Journal*. 2016;15(2): 45-48.
17. Darmstadt GL, Lawn JE, Costello A. Advancing the state of the world's newborns. *Bull World Health Organ*. 2003;2: 224-5.
18. National Institute of Population Research and Training. *Bangladesh demographic and health survey 2014*. Dhaka: National Institute of Population Research and Training. 2014. page 45.
19. Wu XJ, Zhang XD, Shi LP. Retrospective analysis of elective caesarean section and respiratory distress syndrome in the term neonates. *Zhonghua Er Ke Za Zhi*;2009; 47:658-61.
20. Koivisto M, Marttila R, Kurkinen-Raty M, Saarela T, Pokela ML, Jouppila P, et al. Changing incidence and outcome of infants with respiratory distress syndrome in the 1990's: a population-based survey. *Acta Paediatr*. 2004; 11:177-84.
21. Ghafoor T, Mahmud S, Ali S, Dogar SA. Incidence of Respiratory Distress Syndrome. *J Coll Surg Pak*. 13. 2003; 6:271-273
22. Mehrabadi A, Lisonkova S and Joseph K. S. Heterogeneity of respiratory distress syndrome: risk factors and morbidity associated with early and late gestation disease. *BMC Pregnancy and Childbirth* 2016;16: 281-85
23. Luerti M, Parazzini F, Agarossi A, Bianchi C, Rocchetti M, Bevilacqua G. Risk factors for respiratory distress syndrome in the newborn. A multicenter Italian survey. Study Group for Lung Maturity of the Italian Society of Perinatal Medicine. *Acta Obstet Gynecol Scand*.1993; 72: 359-64

24. Desai P.D., Vora H. D. and Mansuri S H. Study of respiratory distress syndrome in preterm neonates. *Indian Journal of Research*. 2014;3(2):218-220.
25. Bryan H, Hawrylyshyn P, Hogg-Johnson S, Inwood S, Finley A, D'Costa M, Chipman M. Perinatal factors associated with the respiratory distress syndrome. *Am J Obstet Gynecol*. 1990;162(2):476-81.
26. Goldberg JD, Cohen WR, Friedman EA. Cesarean section indication and the risk of respiratory distress syndrome. *Obstet Gynecol*. 1981;57(1):30-2
27. Khalid A.Y. Cesarean route of delivery and hyaline membrane disease: a hospital-based case-control study in Greater Beirut. *American University of Beirut, USA*. October. 2003.
28. Liu J., Yang N., and Liu Y. High-risk Factors of Respiratory Distress Syndrome in Term Neonates: A Retrospective Case-control Study. 2014. doi: 10.5152/balkanmedj .2014.8733
29. Bowen JR, Leslie GI, Arnold JD, Jones MP, Gallery ED. Increased incidence of respiratory distress syndrome in infants following pregnancies complicated by hypertension. *Aust N Z J Obstet Gynaecol*. 1988;28(2):109-12.

Comparative Study of neck swelling by clinical, cytological and histopathological examination – a study of 50 cases

Ali MI¹, Huq MM², Rahman SH³, Tarafder KH⁴

Article info

Received : 15.11.2022
Accepted : 14.03.2023
No. of Tables : 08
No. of Figure : 0
No. of References : 25

Cite the Article:

Ali MI, Huq MM, Rahman SH, Tarafder KH. Comparative Study of neck swelling by clinical, cytological and histopathological examination – a study of 50 cases. Pabna Med. Cal. J. 2023;2(1): page 18-23.

Keywords: Close clinicopathological correlation is absolutely necessary for useful clinical interpretation.

Abstract:

Objective: In this study 50 patients of neck swelling were studied to compare the findings with clinical diagnosis and histopathological report for its diagnostic reliability.

Methods: This cross sectional study was done in Department of Otolaryngology – Head & Neck Surgery, Bangabandhu Sheikh Mujib Medical University, Dhaka from July 2015 to June 2017. Fine needle aspiration cytology, MRI, CT scan, and histopathological examinations of postoperative specimen of neck swelling, presentation of analyzed data by various tables, graphs and figures done.

Results: In the present series, 50 cases of neck swelling were studied. Out of these 50 cases 19 were male and 31 were female. The male, female ratio was 1:1.63. Age range was 4 – 60 years with maximum frequency in the 4th decade with the 3rd and 2nd in the following suit. Most of the patient were poor. Clinical, cytological and histopathological diagnosis were available in all the cases. The three sorts of diagnoses were compared with each other. Histopathologically 12 cases were tubercular lymphadenopathy. Metastatic carcinoma and lymphoma, 5 cases for each. 12 were nodular goiter, 5 were thyroid carcinoma. Rest were benign, congenital and nonspecific inflammatory conditions. Correct diagnosis was made by FNAC in 45 cases. In the rest 5 cases, smear were unsatisfactory in 2 cases and gives inconclusive result, remaining 3 were follicular neoplasm and no definitive result were made which were subsequently diagnosed by histopathological examination as a follicular adenoma in 1, and follicular carcinoma in 2 cases. Sensitivity of FNAC in the diagnosis of neck masses were found 91% for tuberculosis, 100% for metastatic carcinoma also for salivary gland tumour. In case of nodular goiter sensitivity was 92%. But it is only 60% sensitive in case of thyroid malignancy, as FNAC cannot demarcate clearly between follicular adenoma and follicular cell carcinoma. But its accuracy in diagnosing papillary cell carcinoma of thyroid was 100%.

Conclusion: Keeping the limitations in mind, FNAC can reduce substantially the need of open biopsy for histopathological examination. The study with the popular saying of Stewart "Diagnosis by aspiration is as reliable as the combined intelligence of the clinician and pathologist makes it".

Introduction:

Swelling in the head and neck region is one of the commonest clinical presentation in the otolaryngologic practice. Presence of a neck mass pose a diagnostic dilemma for the Otolaryngologist. A great number of disease manifest as a palpable and /or

visible swellings in the neck. These may be congenital/developmental, inflammatory/ reactionary or neoplastic (primary/secondary).

Each disease may have different mode of presentation. Conversely many diseases may present with similar

1. Dr. Mohammad Idrish Ali, Consultant, Department of Otolaryngology-Head & Neck Surgery, BSMMU, Dhaka

2. Dr. Md. Mahmudul Huq, Assistant Professor, ENT Department, Khulna Medical College Hospital, Khulna

3. Prof. Sheikh Hasanur Rahman, Professor, Department of Otolaryngology-Head & Neck Surgery, BSMMU, Dhaka

4. Prof. Kamrul Hasan Tarafder, Professor, Department of Otolaryngology-Head & Neck Surgery, BSMMU, Dhaka

Correspondence: Dr. Mohammad Idrish Ali, Consultant, Department of Otolaryngology-Head & Neck Surgery, BSMMU, Mob: 01712-790646, Email: dridrishalient72@gmail.com

symptoms. For this reasons diagnosis often becomes difficult in neck swelling. Many case of neck swelling may be diagnosed after a comprehensive history and a thorough clinical examination of the head neck. Further evaluation is done by hematological, cytological and imaging technique. Diagnosis is confirmed by histopathological examination of the specimen.

The medical profession must be emphatically remained of the frequency with which cervical metastasis may appear as the first and only symptom in cancer of the mouth, pharynx and larynx, less often elsewhere in the body. There can be no possibility of cure until the primary lesion is found. The immediate removal of a lymph node for diagnosis in the does never go best interest of the patient. This procedure should be deferred and used only as a last diagnostic resort.¹

For this reason, the head neck surgeon have advocated a careful search for primary malignancy before the presenting neck lump is biopsied. Open biopsy causes seedling of tumour cell into avascular plane making them resistant to curative radiotherapy or chemotherapy and the placing of a biopsy incision in an area which may subsequently be inappropriate for radical neck dissection flaps^{2,3,4}

Open biopsy is an invasive procedure requiring surgical skill and facilities. It is time consuming, costly and some time hospitalization and general anaesthesia may be needed. It delays the definitive treatment.^{1,3} Fine needle aspiration cytology is a simple procedure that can be done on an outpatient basis without local anaesthetic and gives rapid result. It is simple, cost-effective, less traumatic. The procedure may be repeated several times to obtain adequate material for cytological analysis^{6,7}.

In our country FNAC is gradually becoming more popular as a preoperative highly sensitive and cost effective diagnostic tools.

Through this study, exploration of spectrum of diseases producing neck swelling and their socio-demographic feature done. Histopathological report has been taken as a confirmatory diagnostic test in all the cases and compared with the FNAC findings to find out its diagnostic reliability.

Methods:

This cross sectional study was done in Department of Otolaryngology -Head & Neck Surgery, Bangabandhu Sheikh Mujib Medical University, Dhaka from July 2015 to June 2017. Fine needle aspiration cytology, MRI, CT scan, and histopathological examination of postoperative specimen of neck swelling, presentation of analyzed data by various tables, graphs and figures done.

A total of 50 patients irrespective of age and sex with neck swelling randomly selected for the study who admitted in the Department of Otolaryngology-Head & Neck Surgery, Bangabandhu Sheikh Mujib Medical University Hospital Dhaka from July 2015 to June 2017.

This was a cross sectional study with a sample size 50. Method used in this study included.

Clinical presentation, Fine needle aspirating cytology, Imaging of different types in selective cases, Histopathological examination of post-operative specimen of neck swelling.

All information recorded in a standardized data collection sheet for the study. Data were compiled, analyzed and tabulated in order to obtain statistical and comprehensive results of the study.

Results and observation:

Table-I: Age distribution of the patients.

Age group	No of cases	Percentage
0 – 10	2	4
11 – 20	10	20
21 – 30	14	28
31 – 40	16	32
41 – 50	6	12
51 – 60	2	4
Total	50	100

Table-II: Socioeconomic condition (n=50).

Socioeconomic condition	No of cases	Percentage
Poor	30	60%
Middle class	15	30%
Affluent	5	10%

Table-III: Clinical diagnosis of neck masses.

Clinical diagnosis	Disease	Total No of cases	Percentage (%)
Cervical lymph adenopathy	Tubercular lymph adenitis	19	38
	Metastatic carcinoma	5	10
Thyroid and related neck mass	Goiter	18	36
	Thyroglossal cyst	1	2
Salivary gland swelling	Salivary gland tumour	4	8
Other congenital and developmental neck mass	Branchial cyst	1	2
	Submental dermoid	2	4
	Total	50	100

Table-IV: FNAC diagnosis of neck masses

Clinical diagnosis	FNAC diagnosis	Total No of cases	Percentage (%)
Cervical lymph adenopathy (24)	Tubercular lymph adenitis	10	20
	Metastatic carcinoma	5	10
	Non specific lymphadenitis	3	6
	Lymphoma	5	10
	Unsatisfactory smear	1	2
Thyroid and related neck mass(19)	Nodular goiter	11	22
	Papillary carcinoma of thyroid gland	3	6
	Follicular neoplasm	3	6
	Thyroglossal cyst	1	2
	Unsatisfactory smear	1	2
Salivary gland) swelling (4	Pleomorphic adenoma of salivary gland	3	6
	No specific submandibular sialoadenitis	1	2
Other congenital and developmental neck mass(3)	Branchial cyst	1	2
	Submental dermoid	1	2
	Unsatisfactory smear	1	2

Table-V: Histopathological diagnosis of Neck swelling.

Clinical diagnosis	Histopathological diagnosis	Total No of cases	Percentage (%)
Cervical lymph adenopathy (24)	Tubercular lymph adenitis	11	22
	Metastatic carcinoma	5	10
	Non specific lymphadenitis	3	6
	Lymphoma	5	10
Thyroid and related neck mass(20)	Nodular goiter	12	24
	Papillary carcinoma of thyroid gland	3	6
	Follicular carcinoma	2	4
	Follicular adenoma	1	2
	Thyroglossal cyst	2	4
Salivary gland swelling (4)	Pleomorphic adenoma of salivary gland	3	6
	No specific submandibular sialoadenitis	1	2
Other congenital and developmental neck mass(3)	Branchial cyst	1	2
	Submental dermoid	1	2
	Total	50	100

Table-VI: Comparison between clinical FNAC and histopathological diagnosis of cervical lymph node masses

Clinical diagnosis	No of cases	FNAC diagnosis	No of cases	Histopathological diagnosis	No of cases
Tuberculosis	19	Tubercular lymphadenitis	10	Tubercular lymph adenitis	10
		Non specific lymph adenitis	3	Non specific lymph adenitis	3
		Lymphoma	5	Lymphoma	5
		Unsatisfactory smear	1	Tubercular lymph adenitis	1
Metastatic carcinoma	5	Metastatic carcinoma	5	Metastatic carcinoma	5

Table-VII: Compare between clinical FNAC and histopathological diagnosis of thyroid and related disease

Clinical diagnosis	No of cases	FNAC diagnosis	No of cases	Histopathological diagnosis	No of cases
Goiter	18	Nodular goiter	11	Nodular goiter	11
		Papillary carcinoma of thyroid	3	Papillary carcinoma of thyroid	3
		Follicular neoplasm	3	Follicular carcinoma thyroid	2
		Unsatisfactory smear	1	Follicular adenoma	1
Thyroglossal cyst	1	Thyroglossal cyst	1	Nodular goiter	1
				Thyroglossal cyst	1

Table-VIII: Compare between clinical, FNAC and histopathological diagnosis of salivary gland disease

Clinical diagnosis	No of cases	FNAC diagnosis	No of cases	Histopathological diagnosis	No of cases
Salivary gland tumour	4	Pleomorphic adenoma	3	Pleomorphic adenoma	3
		Chronic submandibular sialoadenitis	1	Chronic submandibular sialoadenitis	1

Discussion

Patients with palpable and or visible neck mass is a quite common presentation to an Otolaryngologist. So one should be rational and methodical for the diagnosis and management of such a patient. In the present study 1 tried to establish the role of FNAC in the management of such patients by establishing its diagnostic sensitivity and specificity in comparison with histopathological one, which is a accurate but cost effective, time consuming and also an invasive procedure.

In the present study of analyzed FNAC report of 50 cases, satisfactory smears were found in 47(94%) cases. In 3(6%) cases smears were unsatisfactory as they showed in adequate material, definitive diagnosis were made by histopathological examination. The rate of unsatisfactory smear in this

study is in close proximity, to that of other studies.^{8,10,11}

In this study out of 50 cases 19(38%) were male and 31(62%) were female. The male to female ratio was 1:1.63. The male to female ratio is consistent with the study of other^{8,11,12,13}.

In the present study age of the patient ranged from 4 to 65 years. The highest number of cases were found in 4th decade. This was followed by 3rd and 2nd decades.

In this present series 40% of neck mass were of thyroid in origin which consistent with other studies.^{3,14}

In case of thyroid swelling out of 20 cases, 12 (60%) cases were proved to be multinodular goiter on histopathology and it is comparable with others.^{10,15}

Here FNAC shows highly sensitivity (91%) and specificity (100%) for nodular goiter. But in case of thyroid malignancy its sensitivity is very low (60%) as it cannot demarcate clearly between follicular carcinoma and follicular adenoma. Although highly sensitive (100%) for papillary carcinoma of thyroid. It is comparable with others studies.^{15,16}

Cervical lymphadenopathy scored the first position (48%) of which tubercular lymphadenitis was 11 cases (46%). This is consistent with the findings of other studies^{17,18}. In this study, the sensitivity (91%) and specificity (100%) of FNAC for diagnosing tubercular lymphadenopathy is high and aligned with other studies^{8,11}.

In this study, only one false negative result was found for tubercular lymphadenopathy by FNAC which was due to inadequate aspirate or observer error.

Metastatic carcinoma was found in about 21% of total cervical lymphadenopathy and sensitivity and specificity for diagnosing such lesion is 100% which was aligned with other studies.^{17,18,20,21,22}

In case of salivary gland lesion which scored in the third position in this series, it is found that FNAC is very useful tool for diagnosis as it a nearly 100% sensitive and specific for such lesion.^{2,3,16}

The overall sensitivity and specificity of FNAC in relation to histopathology is 90% and 100% respectively which are compatible with other study.²

The overall accuracy of FNAC was found in 90% which is similar to that of other studies^{2,3,16,23}.

To obtain maximum benefit from the procedure, close co-operation between the surgeon and pathologist is very important. The role of an experienced cytopathologist is critical for correct diagnosis²⁴. Adequate amount of aspirate from the lesion is essential for accurate diagnosis. Operator must be skilled in performing aspiration. The pathologist must be experienced in cytologic interpretation of the material aspirated. Close clinicopathological correlation is absolutely necessary for useful clinical interpretation.⁷

Conclusion

Keeping the limitations in mind, FNAC can reduce substantially the need of open biopsy for histopathological examination. Actually the diagnosis by aspiration is reliable when it is the teamwork of the clinician and pathologist.

References

1. Martin, H, Romien C. The diagnostic significant of a "Lump in the neck: postgrad Med. NY, 1952; 32: 491-500.
2. Young JEM, Archibald SD, Shier KJ. Needle aspiration cytology biopsy in head and neck mass. Am J of Surgery 1981; 142: 484-489.
3. Raju G, Kakar PK et al. Role of fine needle aspiration biopsy in head and neck tumours. The J of Laryngology and Otology 1988, 102: 248-251.
4. Shaheen OH. The thyroid gland. In: Scott Brown's Otolaryngology, 6th edn, Butter Worth-Heinemann Oxford 1997; p. 5/20/1-17.
5. Sadlar TW: Langman's medical embryology, 8th edn, Baltimore, Williams and Wilkins, 1992, pp. 283-295.
6. Lew WYC. Fine needle aspiration cytology. A personal experience with 800 cases, Singapore Med J. 1987; 28(3): 214-219.
7. Aretz HT, Silverman ML, et al. Fine needle aspiration why it deserve another look. Postgrad. Med 1984; 75: 49-56.
8. Russel RCG, Williams NS, Bulstrode CJK. In: Bailey and Love's short practice of surgery, 24th edn. Arnold, London 2004; p. 710-86.
9. Sattar A, Alam M, Haider A et al. Clinico-pathological study of solitary thyroid nodule. Bangladesh J. of Otolaryngology 2003, 9(1/2): 24-27.
10. Salahuddin M. Correlation among clinical diagnosis, FNAC findings and histopathological reports of neck masses. Dissertation submitted to BCPS in 2002.
11. Hanif MA, Alauddin M. The incidence of tubercular cervical lymphadenitis. Bangladesh J of Otolaryngology 1997, 3(2): 42-46.
12. Siddique A, Mahmud CM, Razeer KAA, Ahmed CN. Presentation of cervical lymphadenopathy to the surgeons. Journal of Surgical Sciences 2005, 9(2): 54-57.
13. Watkinson JC, Gaze MN, Wilson J. A. Benign neck disease. In: Stell and Maran's Head neck surgery, 4th edn Butterworth-Heinemann, Oxford 2000; pp. 181- 182.
14. Islam T. Evaluation of thyroid swelling by clinical cytological and histopathological examination. Dissertation submitted to BCPS, Dhaka, 2003.
15. Mondal A, Roychandhuri B.K. Fine needle aspiration cytology (FNAC) in diagnosis of head &

- neck tumours. *Indian Journal of Otolaryngology and Head Neck*, 1992, 1(4): pp. 176-179.
16. Galib MR, Chowdhury MHR, Hoque MS, Anwar MM, Ahsan S. Cervical lymphadenopathy: A study of hundred and twenty cases. *Sir Salimullah Medical Journal* 2005, 13(2): 59-66.
 17. Chessman AD, Holden HB, Richards AES. The neck. In: Mann CV, Russel RCG, editors, *Bailey and Love's short practice of surgery*. 23rd ed. ELBS with Chapman and Hall, 2000; 256-69: 704-5.
 18. Hag IA, Childozie LC, Reyees FA, Koller SM. Fine needle aspiration of head and neck masses. *Acta Cytol* 2003 May-June; 47(3): 387-392.
 19. Burnand KG, Young AE. *The new Airs companion in surgical studies*, 2nd ed. 1999; 451-7, 980-5.
 20. Alam MS. Clinicopathological study of cervical lymphadenopathy. Dissertation submitted to BCPS in 2004.
 21. Pilotti S, Plasma S, Alasio L, Bartali C, Rilke F. Diagnostic assessment of enlarged superficial lymph node by Fine needle aspiration. *Act Cytol* 1993; 37: 853-866.
 22. Sismanis A, Strong MS, Merriam J. Fine needle aspiration Biopsy diagnosis of neck mass. *Otolaryngology clinic N. America* 1980; 13: 420-429.
 23. Watkinson JC, Gaze MN, Willson January. Metastatic neck disease. In: *Stell and Maran's Head Neck Surgery*, 4th edn; Butterworth-Heinemann, Oxford 2000; p. 197-203.
 24. Hibbert J. Metastatic neck disease. In: *Scott-Browin's Otolaryngology*, 6th edn. Butterworth-Heinemann, Oxford 1997; p. 5/17/1-7.
 25. Evens PR. Tumours of the oropharynx and lymphoma of the head and neck. In: *Scott Brown's Otolaryngology*, 6th edn, Butter Worth-Heinemann Oxford 1997; p. 5/14/5.

Visual Acuity of Children of Darisharifpur Village, Pabna

*Mridha MZEA

Article info

Received : 28.09.2022
Accepted : 14.04.0023
No. of Tables : 0
No. of Figure : 08
No. of References : 10

Cite the Article:

Mridha MZEA. Visual acuity of children of Darisharifpur Village, Pabna. Pabna Med. Cal. J. 2023;2(1): page 24-27.

Keywords: Refractive error, myopia, hypermetropia, rural area, visual acuity.

Abstract

Background: Refractive error in children aged 5-15 years is an important public health problem. Myopia and hypermetropia are leading cause of refractive errors in children in this age group. Both which are visually impairing conditions that can be significantly improved through adequate refractive correction. Population based refractive error studies in children have shows that nearly half of the visual impairment associated with correctable refractive error in this age group in not receiving attention specially those children living in disadvantaged social and economic condition

Methods: This is an observational study. One hundred children of a village Darisharifpur, upazilla- Bera, district Pabna were studied

in Random sampling method. Data were collected from children aged between 9-15 years. Refractive error with other related variables were enclosed with data and and that were analyzed.

Results: 100 children of village with visual problem or symptoms related to visual acuity were examined and prevalence of refractive error in my study 11%. Myopia (72.72%) is more common refractive error my study, myopia 46%, hypermetropia 25%, astigmatism 27%. Majority (63.63%) of refractive error was found in between -0.50D to -1.99D in my study . In my study majority of refractive errors are within 6/9 to 6/18.

Conclusions: Myopia and hypermetropia are leading cause of refractive errors in children in this age group. Both which are visually impairing conditions that can be significantly improved through adequate refractive correction.

Introduction

In Bangladesh, 77% of the population of country lives in rural area. According to Bangladesh national blindness and low vision survey 2002, about 3.3 million adult and 1.3 million children are suffering from refractive errors. Refractive error is second largest cause of visual impairment & blindness at all ages in Bangladesh Refractive error can easily be corrected by spectacles. If refractive error is detected and treated early, the visual loss is reversible in case of children. Despite the huge number of people need spectacles, only 3% of them wire spectacles.

In Bangladesh, ophthalmologists are the main provider of this service but number of ophthalmologists are grossly inadequate and they

work mainly in the large cities. Whereas 77% of population of our country lives in rural area. Refractive service is unavailable in rural area beyond Upazilla level.

Statement of problem:

Blind children due to uncorrected refractive error suffer more than blindness in old aged people and it is a greater socio-economic burden for society. Blindness due to refractive error is one of the obstacles for development of personality, education and career opportunities. Observations on London children aged of 5-8 years (Dustan 1932) shows a peak at +2.30D hypermetropia. In Bangladesh several studies were carried out to determine the refractive error especially in school going children.^{3,5,8}

There are 45 million blind and 135 million visually disabled people in the world. Increasing number of blind is big burden and this will be doubled by 2020, if appropriate measures are not taken. In February 1999 WHO taken global initiative to eliminate the avoidable blindness, “**Vision 2020 the right to sight.**”

The national prevalence survey reveals that number of refractive errors in Bangladesh are 27250 adult and 9925 children per million of population. This means there are estimated 3.3 million adult with refractive error cases with <6/12 visual acuity and 1.3 million children with < 6/18 visual acuity.

Total number of population at the age of 40 years need refractive error correction for near works. Refractive error low vision should be given utmost importance specially for children because delayed intervention can lead them blindness.

Child blindness is relatively complex & demanding areas of works. Though the overall number of blind children is low compared to adult in terms of blindness years it is second only to cataract.

The childhood blindness study in Bangladesh revealed that 31% of blindness was due to problem with lens (cataract) and 27% of blindness was due to problem in the corner (vitamin a deficiency) including Glaucoma (4%) and aphakia (5%) of childhood blindness is thus avoidable (preventable and treatable). This study found that 90% of childhood blindness was developed within first 5 years of life there are estimated 40,000 children are blind in Bangladesh. The national workshop on childhood blindness 2003 recommended that planning for the control of blindness in children should be based on catchments of population of 10 millions rather than 1 million used for planning control of blindness in adult.

Objectives of the study:

The objectives of this study is to determine the prevalence and defferent pattern of refracting error among children of village aged between 9-15 years. However the specific objectives are as follows:

1. To find out the existing problems of refractive error in village children.
2. To asses deferent pattern of refractive errors.
3. To suggest measures to solve the problem.

Rationale of the study:

Refractive error in children age 5-15 years is an important public health problem. Myopia and hypermetropia are leading cause of refractive errors in children in this age group. Both which are visually impairing conditions that can be significantly improved through adequate refractive correction. Population based refractive error studies in children have shows that nearly half of the visual impairment associated with correctable refractive error in this age group in not receiving attention specially those children living in underprivilaged social and economic conditions and such refractive error remains uncorrected and unnecessary visual impairment persists.

In Bangladesh assuming a prevalence of 4 % children aged 5-15 years to have visual acuity of less than 6/18. It is estimated that there are approximate 1.3 million children having refractive error, the large majority of them are correctable.

Methods:

This is an observational study.

Place of study.

Study population : Children age between 9-15 years.

Study period : July 2008 to 30th June 2009

Sample size : 100 children

Data collection and analysis:

By purposive sampling method data were collected from children age between 9-15 years. The questionnaire was prepared containing all information regarding the patient’s chief complaints with duration, refractive condition of family. Socio-economical condition of family, finding related to refractive error collected. visual acuity will be tested with the help of snellen’s E chart followed by Ophthalmoscopy and Ratinoscopy were done data analysed by EPI info 6 programme.

Results

Results are given in tabulated form

Table I: Incidence of refractive error among different sex group

Sex	Total No. (%)	Normal No(%)	Error No(%)
Boys	50 (50%)	44(44%)	6(6%)

Table II: Incidence of refractive error among different occupation

Occupation	Total(%)	Normal(%)	Error(%)
Student	96(96%)	86(86%)	10(10%)
Non student	4(4%)	3(3%)	1(1%)
Total	100(100%)	89(89%)	11(11%)

Table III: Pattern of chief complaints of children

Chief complaints	Number	Percent
Headache	5	45.45%
Defective vision	4	36.36%
Incidental	2	18.18%

Table IV: Pattern of refractive error in different age group

Age group	Number	Percent
9-10 years	4	36.36%
11-13years	5	45.45%
14-15years	2	18.18%

Table V: Pattern of refractive error in one or both eyes

Eye	Number	Percent
One eye	2	18.18%
Both eyes	9	81.81%

Table VI: Severity of refractive error

Visual acuity	Number	Percent
<6/12	4	36.36%
6/12-6/18	3	27.27%
>6/18	4	36.36%

Table VII: Pattern of different types of refractive error

Type of refractive error	Number	Percent
Myopia	8	72.72%
Hypermetropia	2	18.18%
Astigmatism	1	9.09%

Table VIII: Refractive error of children using the eye with highest refractive error

Power	Number	Percent
-0.50D to -1.99D	7	63.63%
-2.00D to -3.99D	2	18.18%
>-3.00D	1	9.09%
+0.50D to +2.00D	1	9.09%

Discussion

100 Village children with visual problem or symptoms related to visual acuity were examined and prevalence of refractive error in my study 11%. Studies carried out R. Mowla Johurul⁸, Sharif and Amanullah¹⁰ showed the figures 8.19% and 6.35% respectively. The difference is due to those studies were done in urban area but my study was done in rural area. Myopia (72.72%) is more common refractive error my study also correlated with other observations of Hussain³, Rahman⁷ and Muqtadir⁵ of our country According to Dutta et al¹, myopia 46%, hypermetropia 25%, astigmatism 27%. In Japan by Muslim et al 1960, a greater prevalence of myopia was reported.

Majority (63.63%) of refractive error was found in between -0.50D to -1.99D in my study which is similar to that of Kragha¹⁹⁸⁷⁴ and Grosvenor 1970². In this study majority of refractive errors are within 6/9 to 6/18.

The efforts of Vision 2020 have resulted in increased funding through governmental and non-governmental sources that are instrumental to the elimination of avoidable causes of blindness. It has been pivotal in gaining and unifying advocacy for eye health at an international and national level. The reduced rates of age-related blindness globally are encouraging of the efforts of Vision 2020. However, there is an overall increase in crude prevalence, and this points to the need for a significant upscaling of efforts. The growing population and demographic shifts will present huge challenges in efforts to minimise vision impairment. To achieve the aims of Vision 2020, there will need to be significant up-scaling of current programmes, increased education, and better reporting of outcomes to better target care. There needs to be continued focus on areas of the world where the magnitude of vision loss is the worst

and the resources available are minimal. There will need to be an increased focus to tackle the burden of NCED that will increase owing to epidemiologic shifts that have already taken place in high-income countries but are also encroaching on low- and middle-income countries. The recent COVID-19 pandemic may have an impact on achieving elimination of avoidable blindness as a public health problem, but it is difficult to say what these may be currently. Undoubtedly, Vision 2020 has led to huge milestones in the way we perceive the importance of eye care and our approaches to providing eye care. However, as can be seen from the most recent data on vision we have a while to go.

Limitation and Recommendation:

Al though children of rural area all over the Bangladesh facing the same problem, the present study cover only one village due to time and other constrains, sample size is 100 which is too small is main limitation of the study.

In spite of these limitation it is hopeful that the study will attract other attention who are interested in depth of issue, although the study not comprehensive it is envisaged to provide some food for thinking about the issue and it enable to attract the attention of the policy makers there might have opportunity to take up necessary steps to strengthen school sight testing program.

References

1. Datta A et al. An epidemiological study of ocular condition among primary school children of Calcutta Corportaion. *Ind.J.Opthal.* 1983;16:40.
2. Grosvenor T. Refraction in Polynesian and European. *AMJ. Optom and Arch. AM Acad. Optom.* 1970;4:673-74.
3. Hussain MA. pattern of refractive error in Bangladesh and their management. A dissertation of FCPS (oph) part-II student at.NIO. 1989. Bangladesh.
4. Kragha IKUK. The distribution of refractive error - Nigeria. *Opthal. Physiol.* 1987;7:241-44.
5. Muktadir AKMA . School sight testing program in hildren. *Trans. Ophtal. Soc. Bang.* 1989;16. 11-14.
6. Nurul Haque DM. Amanullah. Refractive error in children, its effect how to overcome it. *Trans. Opthal. Soc. Bang.* 1994;11: 56-59.
7. Rahman MM. Refractive error in city school of Bangladesh. *Trans opthal. Soc. Bang.* 1989;16:26-30.
8. R. Mowla et al. Refractive error amongst school children and around Rangpur town. *Northern Medical Journal.* 1998;7(1): 24-28.
9. Sorsby A et al. Refraction and its components during the growth of the eye from the age of 3yrs. In medical research council (at Britain) special report series no.301, medical research London, 1961;15: 1-67.
10. Sharif MA. Amanullah prevalence of refractive error in urban School children of Bangladesh. *Trans-Opthal. Soc. Bang.* 1991;18:41-47.

